



# Kongeriget Danmark

Patent application No.: PA 2000 00042

Date of filing: 12 January 2000

Applicants: Novo Nordisk A/S  
Novo Allé  
DK-2880 Bagsværd

Boehringer Ingelheim International GmbH  
D-55216 Ingelheim am Rhein  
Tyskland

This is to certify the correctness of the following information:

The attached photocopy is a true copy of the following information:

The specification, claims and abstract as filed with the application  
on the filing date indicated above.



Patent- og  
Varemærkestyrelsen  
Erhvervsministeriet

TAASTRUP 02 October 2001

  
Karin Schlichting  
Head Clerk

A. Adtadet

12 JAN. 2000

PVS

**TITLE**

Substituted Imidazoles, their Preparation and Use

**FIELD OF THE INVENTION**

5 The present invention relates to novel substituted imidazoles, to the use of these compounds as medicaments, to pharmaceutical compositions comprising the compounds, and to a method of treatment employing these compounds and compositions. The present compounds show a high and selective binding affinity to the histamine H3 receptor indicating a histamine H3 receptor antagonistic or  
10 agonistic activity. As a result, the compounds are useful for the treatment of disorders related to the histamine H3 receptor. More particularly, the present compounds possess a histamine H3 receptor antagonistic activity and are accordingly useful in the treatment of disorders in which a histamine H3 receptor blockade is beneficial.

15

**BACKGROUND OF THE INVENTION**

The existence of the histamine H3 receptor has been known for several years and of current interest for the development of new medicaments (see e.g. Stark, H.; Schlicker, E.; Schunack, W., *Drugs Fut.* **1996**, *21*, 507-520; Leurs, R.; Timmerman, H.; Vollinga, R. C., *Progress in Drug Research* **1995**, *45*, 107-165). Recently, the  
20 histamine H3 receptor has been cloned, cf. Lovenberg, T.W. et al, *Molecular Pharmacology*, **June 1999**, *55*, 1101-1107. The histamine H3 receptor is a presynaptic autoreceptor located in both the central and the peripheral nervous system, the skin and in organs such as the lung, the intestine, probably the spleen and the gastrointestinal tract. The histamine H3 receptor has been demonstrated to regulate the re-  
25 lease of histamine and also of other neurotransmitters such as serotonin and acetylcholine. A histamine H3 receptor antagonist would therefore be expected to increase the release of these neurotransmitters in the brain. A histamine H3 receptor agonist, on the contrary, leads to an inhibition of the biosynthesis and release of  
30 histamine and also of other neurotransmitters such as serotonin and acetylcholine. These findings suggest that histamine H3 receptor agonists and antagonists could

be important mediators of neuronal activity. Accordingly, the histamine H3 receptor is an important target for new therapeutics.

Imidazoles similar to the compounds of the present invention have previously been prepared, and their biological properties have been investigated. Thus, WO 98/29119 relates to tetrahydroimidazopyridine farnesyl-protein inhibitors. JP 06312927 discloses tetrahydroimidazopyridine intermediates for preparing angiotensin II inhibitors. WO 93/17701 discloses tetrahydroimidazopyridine intermediates for preparing endothelin receptor-binding peptides. Klutchko, S., et al., *J. Heterocycl. Chem.*, 28(1), 1991, 97-108 relates to synthesis methods for the preparation of imidazole derivatives. GB 2158440 relates to antiviral compounds. Arcari, G.; Bernardi, L.; Cimaschi, R.; Falconi, G.; Luini, F.; Scarponi, U., *Arzneim. Forsch.*, 34, 11, 1984, 1467-1471, relates to tetrahydroimidazopyridine intermediates for the preparation of imidazopiperidines with anti-ulcer and antisecretory activity, and GB 2028798 relates to tetrahydroimidazopyridine intermediates for the preparation of antiulcer and anticholinergic compounds. However, these references neither disclose nor suggest that the imidazoles may have a histamine H3 receptor antagonistic or agonistic activity.

Furthermore, *Chem. Abstr.*, 87, 201535; Hepp, M.; Schunack, W., *Arch. Pharm. (Weinheim Ger.)*, 313, 9, 1980, 756-762; Vitali et al., *Farmaco Ed. Sci.*, 22, 1967, 821; Habermehl; Ecsy, *Heterocycles*, 5, 1976, 127; Vitali; Bertaccini, *Gazz. Chim. Ital.*, 94, 1964, 296; Emmett, J. C.; Durant, G. J.; Ganellin, C. R.; Roe, A. M.; Turner, J. L., *J. Med. Chem.*, 25, 10, 1982, 1168-1174; Nagarajan, K. et al., *Indian J. Chem. Sect. B*, 15, 1977, 629-634; Casella, L.; Gullotti, M., *J. Am. Chem. Soc.*, 103, 21, 1981, 6338-6347; Piper, I. M.; MacLean, D. B.; Kvarnstrom, I.; Szarek, W. A., *Can. J. Chem.*, 61, 1983, 2721-2728; Williams, R. L.; Neergaard, S., *J. Pharm. Sci.*, 71, 1, 1982, 119-120), DE 2700012, EP 589 665, EP 531 874, WO 92/18115, EP 449 521, US 5,091,390, DE 33 02 125 and DE 33 02 126 disclose imidazopyridine derivatives which are stated to be useful either as intermediates or as therapeutically active substances such as angiotensin II antagonists effective to treat hypertension, peripheral kappa opioid receptor activating substances effective to treat inflammatory pain and N-myristoyl

transferase inhibitors effective as anti-cancer agents. However, these references neither disclose nor suggest that the imidazoles may have a histamine H3 receptor antagonistic or agonistic activity.

- 5 Several publications disclose the preparation and use of histamine H3 agonists and antagonists. Thus, US 4,767,778 (corresponding to EP 214 058), EP 338 939, WO 93/14070, EP 531 219, EP 458 661, EP 197 840, EP 494 010, WO 91/17146, WO 93/12108, WO 93/12107, WO 93/12093, US 5,578,616 (corresponding to WO 95/14007), WO 96/38142, WO 96/38141, WO 95/11894, WO 93/20061,  
10 WO 96/40126, WO 95/06037, WO 92/15567 and WO 94/17058 disclose imidazole derivatives having histamine H3 receptor agonistic or antagonistic activity. However, the structures of these imidazole derivatives are quite different from that of the present compounds. Thus, none of the imidazole derivatives disclosed in these publications have a ring structure fused to the imidazole group such as is the case in the present  
15 compounds.

- In view of the arts interest in histamine H3 receptor agonists and antagonists, novel compounds which trigger the histamine H3 receptor would be a highly desirable contribution to the art. The present invention provides such a contribution to the art  
20 being based on the finding that a specific class of substituted imidazole compounds have a high and specific affinity to the histamine H3 receptor and possess histamine H3 receptor antagonistic activity. Some of these substituted imidazole derivatives are novel per se thereby constituting a further aspect of the invention.

- 25 Due to their histamine H3 receptor antagonistic activity the present compounds are useful in the treatment and/or prevention of a wide range of conditions and disorders in which a blockade of the histamine H3 receptor is beneficial. Thus, the compounds may find use e.g. in the treatment of diseases of the central nervous system, the peripheral nervous system, the cardiovascular system, the pulmonary system, the gas-  
30 trointestinal system and the endocrinological system.

## DEFINITIONS

In the structural formulas given herein and throughout the present specification, the following terms have the indicated meaning:

- 5    The term " $C_{1-6}$ -alkyl" as used herein represent a branched or straight hydrocarbon group having from 1 to 6 carbon atoms. Typical  $C_{1-6}$ -alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, pentyl, isopentyl, hexyl, isohexyl and the like.
- 10   The term " $C_{2-8}$ -alkenyl" as used herein represents a branched or straight hydrocarbon group having from 2 to 8 carbon atoms and at least one double bond. Examples of such groups include, but are not limited to, vinyl, 1-propenyl, 2-propenyl, allyl, iso-propenyl, 1,3-butadienyl, 1-butenyl, 2-butenyl, 1-pentenyl, 2-pentenyl, 1-hexenyl, 2-hexenyl, 1-heptenyl, 2-heptenyl, 1-octenyl, 2-octenyl and the like. In a similar way
- 15   the term " $C_{2-6}$ -alkenyl" represents a branched or straight hydrocarbon group having from 2 to 6 carbon atoms and at least one double bond.

The term " $C_{2-8}$ -alkynyl" as used herein represents a branched or straight hydrocarbon group having from 2 to 8 carbon atoms and at least one triple bond. Examples of

- 20   such groups include, but are not limited to, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 1-pentylnyl, 2-pentylnyl, 1-hexynyl, 2-hexynyl, 1-heptynyl, 2-heptynyl, 1-octynyl, 2-octynyl and the like. In a similar way the term " $C_{2-6}$ -alkynyl" represents a branched or straight hydrocarbon group having from 2 to 6 carbon atoms and at least one triple bond.

25

The term " $C_{1-6}$ -alkoxy" as used herein, alone or in combination, refers to the radical  $-O-C_{1-6}$ -alkyl where  $C_{1-6}$ -alkyl is as defined above. Representative examples are methoxy, ethoxy, n-propoxy, isopropoxy, butoxy, *sec*-butoxy, *tert*-butoxy, pentoxy, isopentoxy, hexoxy, isohexoxy and the like.

30

The term "C<sub>1-6</sub>-alkylthio" as used herein, alone or in combination, refers to the radical -S-C<sub>1-6</sub>-alkyl where C<sub>1-6</sub>-alkyl is as defined above. Representative examples are methylthio, ethylthio, isopropylthio, propylthio, butylthio, pentylthio and the like.

- 5 The term "C<sub>3-15</sub>-cycloalkyl" as used herein represents a carbocyclic group having from 3 to 15 carbon atoms such as from 3 to 8 carbon atoms. Representative examples are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl and the like. In the same way the term "C<sub>3-8</sub>-cycloalkyl" represents a carbocyclic group having from 3 to 8 carbon atoms

10

The term "aryl" as used herein is intended to include carbocyclic aromatic ring systems such as phenyl, naphthyl (1-naphthyl or 2-naphthyl), anthracenyl (1-anthracenyl, 2-anthracenyl, 3-anthracenyl), phenanthrenyl, fluorenyl, indenyl and the like. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 1-(1,2,3,4-tetrahydronaphthyl) and 2-(1,2,3,4-tetrahydronaphthyl).

15

The term "aroyl" as used herein refers to the radical -CO-aryl where aryl is as defined above. Non-limiting examples are benzoyl, naphthoyl, anthracenoyl, phenanthrenoyl, fluorenyl, indenoyl and the like.

20

The term "aryloxy" as used herein refers to the radical -O-aryl where aryl is as defined above. Non-limiting examples are phenoxy, naphthoxy, anthracenyloxy, phenantrenyloxy, fluorenyloxy, indenyloxy and the like.

25

The term "arylthio" as used herein refers to the radical -S-aryl where aryl is as defined above. Non-limiting examples are phenylthio, naphthylthio, phenanthrenylthio, fluorenylthio, indenylthio and the like.

The term "arylamino" as used herein refers to the radical -NH-aryl where aryl is as defined above. Non-limiting examples are phenylamino, naphthylamino, phenanthrenylamino, fluorenylamino, indenylamino and the like.

- 5 The term "arylsulfonyl" as used herein refers to the radical -S(=O)<sub>2</sub>-aryl where aryl is as defined above. Non-limiting examples are phenylsulfonyl, naphthylsulfonyl, phenanthrenylsulfonyl, fluorenylsulfonyl, indenylsulfonyl and the like.

The term "heteroaryl" as used herein is intended to include heterocyclic aromatic ring systems containing one or more heteroatoms selected from nitrogen, oxygen and sulfur such as furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyranyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, tetrazolyl, thiadiazinyl, indolyl, isoindolyl, benzofuryl, benzothienyl, benzothiophenyl (thianaphthenyl), indazolyl, benzimidazolyl, benzthiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, purinyl, quinazolinyl, quinoliziny, quinoliny, isoquinoliny, quinoxaliny, naphthyridiny, pteridinyl, carbazolyl, azepiny, diazepiny, acridiny and the like. Heteroaryl is also intended to include the partially or fully hydrogenated derivatives of the heterocyclic systems enumerated above. Non-limiting examples of such partially or fully hydrogenated derivatives are pyrroliny, pyrazoliny, indoliny, pyrrolidinyl, piperidinyl, piperazinyl, azepiny, diazepiny, morpholiny, thiomorpholiny, oxazolidiny, oxazoliny, oxazepiny, aziridiny and tetrahydrofurany.

25

The term "heteroaroyl" as used herein refers to the radical -CO-heteroaryl where heteroaryl is as defined above. Non-limiting examples are furoyl, thienylcarbonyl, pyridoyl, oxazolylcarbonyl, benzofurylcarbonyl, benzimidazolylcarbonyl, pyrroliny, azepiny, diazepiny, morpholiny, thiomorpholiny, oxazolidiny, oxazoliny, oxazepiny, aziridiny and tetrahydrofurany.

30

The term "heteroaryloxy" as used herein refers to the radical -O-heteroaryl where heteroaryl is as defined above. Non-limiting examples are furyloxy, thienyloxy, pyridyloxy, oxazolyloxy, benzofuryloxy, benzimidazolyloxy, pyrrolinyloxy, azepinyloxy and the like.

5

The term "heteroarylamino" as used herein refers to the radical -NH-heteroaryl where heteroaryl is as defined above. Non-limiting examples are furylamino, thienylamino, pyridylamino, oxazolylamino, benzofurylamino, benzimidazolylamino, pyrrolinylamino, azepinylamino and the like.

10

The term "heteroarylthio" as used herein refers to the radical -S-heteroaryl where heteroaryl is as defined above. Non-limiting examples are furylthio, thienylthio, pyridylthio, oxazolylthio, benzofurylthio, benzimidazolylthio, pyrrolinylthio, azepinylthio and the like.

15

The term "heteroarylsulfonyl" as used herein refers to the radical -S(=O)<sub>2</sub>-heteroaryl where heteroaryl is as defined above. Non-limiting examples are furylsulfonyl, thienylsulfonyl, pyridylsulfonyl, oxazolylsulfonyl, benzofurylsulfonyl, benzimidazolylsulfonyl, pyrrolinylsulfonyl, azepinylsulfonyl and the like.

20

The term "acylamino" as used herein represents a radical of the form -N(L)-C(=O)-G where G and L independently represent hydrogen, C<sub>1-6</sub>-alkyl, aryl or heteroaryl as defined above. Non-limiting examples are acetylamino, propanoylamino, butyrylamino, pentanoylamino, benzoylamino, furoylamino, pyridoylamino and the like.

25

The term "sulfonylamino" as used herein represents a radical of the form -N(L)-S(=O)<sub>2</sub>-G where G and L independently represent hydrogen, C<sub>1-6</sub>-alkyl, aryl or heteroaryl as defined above. Non-limiting examples are methanesulfonylamino, propanesulfonylamino, benzenesulfonylamino, N-methyl-N-(benzenesulfonyl)amino, 4-methylbenzenesulfonylamino N-butyl-N-(4-methylbenzenesulfonyl)amino, 2-thienylsulfonylamino and the like.

30




The term "halogen" means fluorine, chlorine, bromine or iodine.

- As used herein, the phrase "3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring" is intended to include carbocyclic rings which are saturated or contain one or more double bonds as well as heterocyclic rings containing one or more heteroatoms selected from nitrogen, oxygen or sulfur which are saturated or contain one or more double bonds.
- 10 The term "optionally substituted" as used herein means that the groups in question are either unsubstituted or substituted with one or more of the substituents specified. When the groups in question are substituted with more than one substituent the substituents may be the same or different.
- 15 As used herein, the phrase "a functional group which can be converted to hydrogen *in vivo*" is intended to include any group which upon administering the present compounds to the subjects in need thereof can be converted to hydrogen e.g. enzymatically or by the acidic environment in the stomach. Non-limiting examples of such groups are acyl, carbamoyl, monoalkylated carbamoyl, dialkylated carbamoyl, alkoxycarbonyl, alkoxyalkyl groups and the like such as C<sub>1-6</sub>-alkanoyl, aroyl, C<sub>1-6</sub>-alkylcarbamoyl, di-C<sub>1-6</sub>-alkylcarbamoyl, C<sub>1-6</sub>-alkoxycarbonyl and C<sub>1-6</sub>-alkoxy-C<sub>1-6</sub>-alkyl.
- 20

- Certain of the above defined terms may occur more than once in the structural formulas, and upon such occurrence each term shall be defined independently of the other.
- 25

## DESCRIPTION OF THE INVENTION

The present invention relates to novel, substituted imidazoles of the general formula I



5

wherein

R<sup>1</sup> is hydrogen or a functional group which can be converted to hydrogen *in vivo*;

10 R<sup>2</sup> is hydrogen, C<sub>1-6</sub>-alkyl, halogen, cyano, trifluoromethyl, hydroxy or -NR<sup>7</sup>R<sup>8</sup>,

wherein R<sup>7</sup> and R<sup>8</sup> independently are

hydrogen,

15

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

20

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroaryl-amino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

5 C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

10 R<sup>7</sup> and R<sup>8</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> independently are

hydrogen, carboxy, C<sub>1-6</sub>-alkoxycarbonyl, cyano, trifluoromethyl, halogen,

20

C<sub>3-8</sub>-cycloalkyl optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

25

C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with

C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, cyano, halogen, trifluoromethyl, carboxy, C<sub>1-6</sub>-alkoxycarbonyl,

30

C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

5 aryl optionally substituted with

halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, heteroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

10 -CO-NR<sup>9</sup>R<sup>10</sup>,

wherein R<sup>9</sup> and R<sup>10</sup> independently are

hydrogen,

15

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

20

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

25

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

5 R<sup>9</sup> and R<sup>10</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl,  
10 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, and/or R<sup>5</sup> and R<sup>6</sup> together with the carbon atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;  
15

20 m, n, p and q independently are 0, 1 or 2;

X is a valence bond, -CH<sub>2</sub>-, -C(=O)-, -C(=S)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)-, -C[=C(CN)<sub>2</sub>]-, -C(=CH-CN)-, -C(=NR<sup>11</sup>)- or -C(=N-S(=O)<sub>2</sub>R<sup>11a</sup>)-,

25 wherein R<sup>11</sup> is

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with  
30 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

5 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

10

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

15

R<sup>11a</sup> is

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

20

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

25

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

30

Y is a valence bond, -O- or -N(R<sup>12</sup>)-,

wherein R<sup>12</sup> is

hydrogen,

5

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

10

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroaryl-amino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

15

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

20

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

A is a valence bond, C<sub>1-8</sub>-alkylene, C<sub>2-8</sub>-alkenylene, C<sub>2-8</sub>-alkynylene, C<sub>3-8</sub>-cyclo-alkylene or phenylene, or

25

when Y is -N(R<sup>12</sup>)-, A, together with R<sup>12</sup> and the nitrogen atom to which they are connected, may form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring system optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl,

30

heteroaryl, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and

Z is -R<sup>13</sup>, -OR<sup>13</sup>, -SR<sup>13</sup>, -NR<sup>13</sup>R<sup>14</sup>, -CHR<sup>13</sup>R<sup>14</sup>, -CR<sup>13</sup>R<sup>14</sup>R<sup>15</sup> or =CR<sup>13</sup>R<sup>14</sup>,

5

wherein R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> independently are

hydrogen,

10

C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with aryl, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, heteroaryl or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

15

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino, halogen, cyano or trifluoromethyl,

20

aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, C<sub>3-8</sub>-cycloalkanecarbonyl, hydroxy,

25

amino, halogen, cyano or trifluoromethyl, where

30

R<sup>13</sup> and R<sup>14</sup> or R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup>, when they do not represent hydrogen, may be joined by one or more bridging linkers such as a valence bond, C<sub>1-4</sub>-alkylene, C<sub>2-4</sub>-alkenylene, -O-, -S-, -N(R<sup>16</sup>)-, -C(=O)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(R<sup>16</sup>R<sup>17</sup>)-, phenylene, biphenylene, -O-C<sub>1-4</sub>-alkylene, -S-C<sub>1-4</sub>-alkylene, -N(R<sup>16</sup>)-C<sub>1-4</sub>-alkylene,



-N=C<sub>1-4</sub>-alkylene, -O-C<sub>2-4</sub>-alkenylene, -S-C<sub>2-4</sub>-alkenylene or -N(R<sup>16</sup>)-C<sub>2-4</sub>-alkenylene, to form a mono-, bi- or polycyclic ring system,

wherein R<sup>16</sup> and R<sup>17</sup> independently are

5

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

10

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, hetero-  
aryl amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen,  
cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, aryl-  
sulfonyl, arylamino or heteroaryl amino,

15

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which  
are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-  
amino or heteroaryl amino,

20

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen,  
cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsul-  
fonyl, arylamino or heteroaryl amino; or

25

R<sup>16</sup> and R<sup>17</sup>, together with the nitrogen atom to which they are connected,  
form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring op-  
tionally containing one or more further heteroatoms selected from nitro-  
gen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl,

30

C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

with the provisos that

5

when X is -CS-,  $R^1 = R^2 = R^5 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ , the group -Y-A-Z must not start with the radical -NH-;

10

when the group -X-Y-A-Z starts with the radical -CH<sub>2</sub>-,  $R^1 = R^2 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy or aminocarbonyl;

15

when X is -CO-, the group -Y-A-Z starts with the radical -NH-,  $R^1 = R^2 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ , the remainder of the group -Y-A-Z must not be hydrogen, unsubstituted or C<sub>1-6</sub>-alkoxy substituted phenyl, unsubstituted C<sub>3-8</sub>-cycloalkyl or unsubstituted C<sub>1-6</sub>-alkyl;

20

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl,  $R^1 = R^2 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy, aminocarbonyl or 4-phenyl-piperazin-1-ylcarbonyl;

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl,  $R^1 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $R^2 = \text{butyl}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be methoxycarbonyl;

25

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl,  $R^1 = R^2 = R^4 = R^5 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^3$  must not be hydrogen, ethyl, isopropyl or phenyl;

when X is -CO-, Y is -O-, A is a valence bond, Z is *tert*-butyl,  $R^1 = R^2 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy;

30

when X is -CO-, Y is -O-, A is a valence bond, Z is *tert*-butyl,  $R^1 = R^2 = R^4 = R^5 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^3$  must not be 4-cyanophenyl;

when X is -CO-, the group -Y-A-Z starts with the radical -O-,  $R^1 = R^2 = R^6 =$  hydrogen,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy, aminocarbonyl or hydrogen;

- 5 when -X is -CO-, the group -Y-A-Z starts with the radical -CH<,  $R^1 = R^2 = R^3 = R^4 = R^6 =$  hydrogen,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be hydroxymethyl,  $C_{1-6}$ -alkoxy-carbonyl or carboxy; and

- 10 when X is -CO-, the group -Y-A-Z is 4-methoxyphenyl,  $R^1 = R^2 = R^3 = R^4 = R^6 =$  hydrogen,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy;

as well as any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof.

- 15 Preferably, m, n, and p are 0 and q is 1.

Preferably,  $R^1$  and  $R^2$  are both hydrogen.

Preferably,  $R^5$  and  $R^6$  are both hydrogen.

20

In a preferred embodiment of the invention m, n, and p are 0 and q is 1;  $R^1$  and  $R^2$  are hydrogen;  $R^5$  and  $R^6$  are hydrogen; X is -C(=O)-; Y is a valence bond; A is a valence bond or  $C_{1-8}$ -alkylene; and Z is  $-R^{13}$ ,  $-CHR^{13}R^{14}$  or  $-NHR^{13}R^{14}$ , wherein  $R^{13}$  and  $R^{14}$  are as defined above.

25

When Z is  $-R^{13}$ ,  $R^{13}$  is preferably

aryl,  $C_{3-15}$ -cycloalkyl, aroyl or heteroaryl, which are optionally substituted with  
 aryl- $C_{1-6}$ -alkyl, heteroaryl- $C_{1-6}$ -alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-  
 30 amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl,  $C_{1-6}$ -alkylsulfonyl,

sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl.

When Z is -CHR<sup>13</sup>R<sup>14</sup>, R<sup>13</sup> and R<sup>14</sup> are preferably independently

5

hydrogen, or

aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with

10

aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl.

When Z is -NR<sup>13</sup>R<sup>14</sup>, R<sup>13</sup> and R<sup>14</sup> are preferably independently

15

hydrogen,

aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with

20

aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl, or

R<sup>13</sup> and R<sup>14</sup> are each independently aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which

25

are joined with a C<sub>1-4</sub>-alkylene group to form a polycyclic ring system.

R<sup>3</sup> and R<sup>4</sup> are preferably independently

hydrogen,

30

C<sub>3-8</sub>-cycloalkyl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

5 C<sub>1-6</sub>-alkyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino  
10 or heteroarylamino,


aryl optionally substituted with

halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, heteroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or  
15 -CO-NR<sup>9</sup>R<sup>10</sup> wherein R<sup>9</sup> and R<sup>10</sup> are as defined above for formula I, or

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, form a C<sub>3-8</sub>-cycloalkyl ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino.  
20

More preferably, R<sup>3</sup> and R<sup>4</sup> are both hydrogen or are both C<sub>1-6</sub>-alkyl such as methyl or ethyl or R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, form a C<sub>3-8</sub>-cycloalkyl ring such as spirocyclobutane or spirocyclopentane.  
25

Specific examples of the above preferred embodiments of the present invention are the following compounds:



5

wherein

Ex. No.	R <sup>3</sup>	R <sup>4</sup>	X	Y	A	Z
14-002	H	H	CO	bond	methylene	cyclohexyl
14-004	H	H	CO	bond	ethylene	4-fluorobenzoyl
14-003	H	H	CO	bond	bond	cyclohexyl
14-005	H	H	CO	bond	pentylene	benzoyl
4	H	H	CO	bond	ethylene	cyclohexyl
3	H	H	CO	bond	pentylene	phenyl
2	H	H	CO	bond	butylene	cyclohexyl
	H	H	CO	bond	bond	methyl
6	H	H	CO	bond	methylene	diphenylmethyl
8	H	H	CO	bond	ethylene	10,11-dihydro-5H-dibenzo[b,f]azepin-5-yl
1	CH <sub>3</sub>	CH <sub>3</sub>	CO	bond	ethylene	cyclohexyl
5	H	4-isopropylphenyl	CO	bond	ethylene	4-fluorophenyl
7	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>3</sub>	CO	bond	ethylene	cyclohexyl
9	spiro-	-(CH <sub>2</sub> ) <sub>3</sub> -	CO	bond	ethylene	cyclohexyl

- 10 In another preferred embodiment m, n, and p are 0 and q is 1; R<sup>1</sup> and R<sup>2</sup> are hydrogen; R<sup>5</sup> and R<sup>6</sup>; X is -C(=O)-; Y is -NH-; A is a valence bond, C<sub>1-8</sub>-alkylene or C<sub>3-8</sub>-cycloalkylene; and

Z is -R<sup>13</sup> in which R<sup>13</sup> is

5390.029-DK,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, heteroaryl

5 or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroaryl

arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryl-

oxy, acylamino, hydroxy, amino, halogen, cyano or trifluoromethyl,

10

aryl, C<sub>3-15</sub>-cycloalkyl or heteroaryl, which are optionally substituted with

aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-

amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl,

sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy,

15 C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl, or

Z is -CHR<sup>13</sup>R<sup>14</sup> in which R<sup>13</sup> and R<sup>14</sup> independently are

hydrogen,

20

C<sub>1-6</sub>-alkyl optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro,

arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl,

C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hy-

25 droxy, amino, halogen, cyano or trifluoromethyl,

aryl, C<sub>3-15</sub>-cycloalkyl or heteroaryl, which are optionally substituted with

aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-

amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl,


30 sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy,

C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl.

R<sup>3</sup> and R<sup>4</sup> are preferably as defined above in the preferred embodiment.

More preferably, R<sup>3</sup> and R<sup>4</sup> both represent hydrogen, or one of R<sup>3</sup> and R<sup>4</sup> represents  
 5 hydrogen while the other represents C<sub>3-8</sub>-cycloalkyl substituted C<sub>1-6</sub>-alkyl.

Specific examples of the above preferred embodiments of the present invention are the following compounds:



10

wherein

Ex. No.	R <sup>3</sup>	R <sup>4</sup>	X	Y	A	Z
10-001	H	H	CO	NH	-(CH <sub>2</sub> ) <sub>2</sub> -	2-thienyl
10-002	H	H	CO	NH	bond	3,5-dimethyl-1,2-oxazol-4-yl
10-003	H	H	CO	NH	-CH(CH <sub>3</sub> )-	1-naphthyl
10-004	H	H	CO	NH	bond	2-phenylcyclopropyl
10-005	H	H	CO	NH	bond	1-(4-bromophenyl)ethyl
10-006	H	H	CO	NH	bond	2-(trifluoromethyl)phenyl
10-007	H	H	CO	NH	ethylene	phenyl
10-008	H	H	CO	NH	bond	4-(trifluoromethyl)phenyl
10-009	H	H	CO	NH	bond	3-cyanophenyl
10-010	H	H	CO	NH	bond	4-cyanophenyl
10-011	H	H	CO	NH	bond	n-octyl
11	H	H	CO	NH	methylene	2,4-dichlorophenyl
12	H	2-cyclohexylethyl	CO	NH	bond	ethyl
13	H	2-cyclohexylethyl	CO	NH	methylene	2,4-dichlorophenyl



In a further preferred embodiment of the invention  $m = n = p = 0$  and  $q = 1$ ;  $R^1 = R^2 =$  hydrogen;  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  are hydrogen; X is -C(O)- and Y is -O-.

5 Preferably, A is  $C_{1-6}$ -alkylene or a valence bond.

More preferably, A is methylene or ethylene.

10 Z is preferably  $-R^{13}$ ,  $-\text{CHR}^{13}\text{R}^{14}$  or  $-\text{CR}^{13}\text{R}^{14}\text{R}^{15}$  wherein  $R^{13}$ ,  $R^{14}$  and  $R^{15}$  are as defined above for formula I.

When Z is  $-R^{13}$ ,  $R^{13}$  is preferably aryl, heteroaryl or  $C_{3-15}$ -cycloalkyl which may optionally be substituted as defined above for formula I.

15 More preferably, Z is phenyl, naphthyl, thienyl, cyclopentyl, cyclohexyl or cyclohexenyl which may optionally be substituted as defined above for formula I.

In one preferred embodiment thereof Z is phenyl, naphthyl, thienyl, cyclopentyl, cyclohexyl or cyclohexenyl.


20

In another preferred embodiment thereof Z is phenyl which is substituted with one to three substituents selected from  $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxy, halogen, phenyl and trifluoromethyl.

25 When Z is  $-\text{CHR}^{13}\text{R}^{14}$ ,  $R^{13}$  is preferably  $C_{1-6}$ -alkyl and  $R^{14}$  is preferably phenyl.

When Z is  $-\text{CR}^{13}\text{R}^{14}\text{R}^{15}$ ,  $R^{13}\text{R}^{14}$  and  $R^{15}$  are preferably each  $C_{1-6}$ -alkyl which are joined with  $C_{1-4}$ -alkylene linkers to form a polycarbocyclic ring system, such as adamantyl.

















Specific examples of the above preferred embodiments of the present invention are the following compounds:







5

wherein

Ex. No.	R <sup>3</sup>	A	Z
15	H	-CH <sub>2</sub> -	
16	H	-CH <sub>2</sub> -	
17	H	-CH <sub>2</sub> -	
18	H	-CH <sub>2</sub> -	
19	H	-CH <sub>2</sub> -	
20	H	-CH <sub>2</sub> -	
21	H	-CH <sub>2</sub> -	
22	H	-CH <sub>2</sub> -	
23	H	-(CH <sub>2</sub> ) <sub>2</sub> -	
24	H	-(CH <sub>2</sub> ) <sub>2</sub> -	

25	H	-CH <sub>2</sub> -	
26	H	-CH <sub>2</sub> -	
27	H	-CH <sub>2</sub> -	
28	H	-CH <sub>2</sub> -	
29	H	-(CH <sub>2</sub> ) <sub>2</sub> -	
30	H	-CH <sub>2</sub> -	
31	H	-CH <sub>2</sub> -	
32	H	-CH <sub>2</sub> -	
33	H	-CH <sub>2</sub> -	
34	H	-CH <sub>2</sub> -	
35	H	-CH <sub>2</sub> -	
36	H	-(CH <sub>2</sub> ) <sub>2</sub> -	
37	H	-(CH <sub>2</sub> ) <sub>2</sub> -	
38	H	-CH <sub>2</sub> -	
39	-CF <sub>3</sub>	-CH <sub>2</sub> -	
40	H	-(CH <sub>2</sub> ) <sub>2</sub> -	

41	H	-CH <sub>2</sub> -	
42	H	-CH <sub>2</sub> -	
43	H	-CH <sub>2</sub> -	
44	H	valence bond	

In another embodiment of the invention  $m = n = p = 0$  and  $q = 1$ ;  $R^1 = R^2 = \text{hydrogen}$ ;  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  are hydrogen; X is -C(O)-; Y is -N(R<sup>12</sup>)- wherein R<sup>12</sup> and A, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring system optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is hydrogen.

In yet another embodiment of the invention  $m = n = p = 0$  and  $q = 1$ ;  $R^1 = R^2 = \text{hydrogen}$ ;  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  are hydrogen; X is -C(O)-; Y is -N(R<sup>12</sup>)- wherein R<sup>12</sup> is C<sub>1-6</sub>-alkyl; A is C<sub>1-8</sub>-alkylene; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is aryl optionally substituted as defined above for formula I.

In still another embodiment of the invention  $m = n = p = 0$  and  $q = 1$ ;  $R^1 = R^2 = \text{hydrogen}$ ;  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  are hydrogen; X is -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)- or -C(=N-S(=O)<sub>2</sub>R<sup>11a</sup>)- wherein R<sup>11a</sup> is C<sub>1-6</sub>-alkyl or phenyl substituted with C<sub>1-6</sub>-alkyl; Y is -NH-; A is C<sub>1-8</sub>-alkylene; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is aryl or C<sub>3-15</sub>-cycloalkyl optionally substituted as defined for formula I.

R<sup>13</sup> is preferably phenyl or cyclohexyl which are optionally substituted with halogen, C<sub>1-6</sub>-alkyl or C<sub>3-8</sub>-cycloalkanecarbonyl.

A is preferably methylene, ethylene or propylene.

In a further embodiment of the invention  $m = n = p = 0$  and  $q = 1$ ;  $R^1 = R^2 =$  hydrogen;  $R^3, R^4, R^5$  and  $R^6$  are hydrogen; X is  $-S(=O)_2-$ ; Y is a valence bond; A is a valence bond or  $C_{1-8}$ -alkylene; and Z is  $-R^{13}$  wherein  $R^{13}$  is aryl or  $C_{3-15}$ -cycloalkyl optionally substituted as defined for formula I.

$R^{13}$  is preferably phenyl or cyclohexyl which are optionally substituted with halogen,  $C_{1-6}$ -alkyl or  $C_{3-8}$ -cycloalkanecarbonyl.


A is preferably methylene, ethylene or propylene.

In still a further embodiment of the invention  $m = n = p = 0$  and  $q = 1$ ;  $R^1 = R^2 =$  hydrogen;  $R^3, R^4, R^5$  and  $R^6$  are hydrogen; X is  $-C(=O)-$ ; Y is  $-NH-$ ; A is a valence bond or  $C_{1-8}$ -alkylene; and Z is  $-R^{13}$  wherein  $R^{13}$  is aryl or  $C_{3-15}$ -cycloalkyl optionally substituted as defined for formula I.

$R^{13}$  is preferably phenyl or cyclohexyl which are optionally substituted with halogen,  $C_{1-6}$ -alkyl or  $C_{3-8}$ -cycloalkanecarbonyl.

A is preferably methylene, ethylene or propylene.

In a further aspect the invention relates to novel, substituted imidazoles of the general formula I''



5 wherein

R<sup>1</sup> is hydrogen or a functional group which can be converted to hydrogen *in vivo*;

R<sup>2</sup> is hydrogen, C<sub>1-6</sub>-alkyl, halogen, cyano, trifluoromethyl, hydroxy or -NR<sup>7</sup>R<sup>8</sup>,

10

wherein R<sup>7</sup> and R<sup>8</sup> independently are

hydrogen,

15

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroaryl-amino,

20

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

5 C<sub>1-6</sub>-alkylsulfonyl optionally substituted with  
C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

10 R<sup>7</sup> and R<sup>8</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;  
15

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> independently are

hydrogen, carboxy, C<sub>1-6</sub>-alkoxycarbonyl, cyano, trifluoromethyl, halogen,

20 C<sub>3-8</sub>-cycloalkyl optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

25 C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with

C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, cyano, halogen, trifluoromethyl, carboxy, C<sub>1-6</sub>-alkoxycarbonyl,

30 C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

5 aryl optionally substituted with

halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, heteroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

10 -CO-NR<sup>9</sup>R<sup>10</sup>,

wherein R<sup>9</sup> and R<sup>10</sup> independently are

hydrogen,

15

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

20 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

25 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with



C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

5 R<sup>9</sup> and R<sup>10</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, 10 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, and/or R<sup>5</sup> and R<sup>6</sup> together with the carbon atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, 15 trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

20 m, n, p and q independently are 0, 1 or 2;

X is a valence bond, -CH<sub>2</sub>-, -C(=O)-, -C(=S)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)-, -C[=C(CN)<sub>2</sub>]-, -C(=CH-CN)- or -C(=NR<sup>11</sup>)-,

25 wherein R<sup>11</sup> is

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with 30 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

5 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

10

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

15

Y is a valence bond, -O- or -N(R<sup>12</sup>)-,

wherein R<sup>12</sup> is

20 hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

25

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

30

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

5 C<sub>1-6</sub>-alkylsulfonyl optionally substituted with  
C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

10 A is a valence bond, C<sub>1-8</sub>-alkylene, C<sub>2-8</sub>-alkenylene, C<sub>2-8</sub>-alkynylene, C<sub>3-8</sub>-cycloalkylene or phenylene, or

when Y is -N(R<sup>12</sup>)-, A, together with R<sup>12</sup> and the nitrogen atom to which they are connected, may form a 3 to 8 membered, saturated or unsaturated, heterocyclic  
15 ring system optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and

20 Z is -R<sup>13</sup>, -OR<sup>13</sup>, -SR<sup>13</sup>, -NR<sup>13</sup>R<sup>14</sup>, -CHR<sup>13</sup>R<sup>14</sup>, -CR<sup>13</sup>R<sup>14</sup>R<sup>15</sup> or =CR<sup>13</sup>R<sup>14</sup>,

wherein R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> independently are

25 hydrogen,

C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with aryl, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hetero-  
30 aryl or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, hetero-  
 aryl-C<sub>1-6</sub>-alkyl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl,  
 arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino,  
 arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino, halogen,  
 cyano or trifluoromethyl,

aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with  
 aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, het-  
 eroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkyl-  
 sulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino,  
 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or  
 trifluoromethyl, where

R<sup>13</sup> and R<sup>14</sup> or R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup>, when they do not represent hydrogen, may be  
 joined by one or more bridging linkers such as a valence bond, C<sub>1-4</sub>-alkylene,  
 C<sub>2-4</sub>-alkenylene, -O-, -S-, -N(R<sup>16</sup>)-, -C(=O)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(R<sup>16</sup>R<sup>17</sup>)-, phen-  
 ylene, biphenylene, -O-C<sub>1-4</sub>-alkylene, -S-C<sub>1-4</sub>-alkylene, -N(R<sup>16</sup>)-C<sub>1-4</sub>-alkylene,  
 -N=C<sub>1-4</sub>-alkylene, -O-C<sub>2-4</sub>-alkenylene, -S-C<sub>2-4</sub>-alkenylene or -N(R<sup>16</sup>)-C<sub>2-4</sub>-al-  
 kenylene, to form a mono-, bi- or polycyclic ring system,

wherein R<sup>16</sup> and R<sup>17</sup> independently are

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, hetero-  
 arylamino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with  
 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen,  
 cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, aryl-  
 sulfonyl, arylamino or heteroaryl-amino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsul-fonyl, arylamino or heteroarylamino, or

R<sup>16</sup> and R<sup>17</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitro-gen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

with the provisos that

when X is -CS-, R<sup>1</sup> = R<sup>2</sup> = R<sup>5</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, the group -Y-A-Z must not start with the radical -NH-;

when the group -X-Y-A-Z starts with the radical -CH<sub>2</sub>-, R<sup>1</sup> = R<sup>2</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>5</sup> must not be carboxy or aminocarbonyl;

when X is -CO-, the group -Y-A-Z starts with the radical -NH-, R<sup>1</sup> = R<sup>2</sup> = R<sup>6</sup> = hydro-gen, m = n = p = 0 and q = 1, the remainder of the group -Y-A-Z must not be hydro-gen, unsubstituted or C<sub>1-6</sub>-alkoxy substituted phenyl, unsubstituted C<sub>3-8</sub>-cycloalkyl or unsubstituted C<sub>1-6</sub>-alkyl;

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl, R<sup>1</sup> = R<sup>2</sup> = R<sup>3</sup> = R<sup>4</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>5</sup> must not be carboxy, aminocarbonyl or 4-phenyl-piperazin-1-ylcarbonyl;

5

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl, R<sup>1</sup> = R<sup>3</sup> = R<sup>4</sup> = R<sup>6</sup> = hydrogen, R<sup>2</sup> = butyl, m = n = p = 0 and q = 1, R<sup>5</sup> must not be methoxycarbonyl;

10

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl, R<sup>1</sup> = R<sup>2</sup> = R<sup>4</sup> = R<sup>5</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>3</sup> must not be hydrogen, ethyl, isopropyl or phenyl;

when X is -CO-, Y is -O-, A is a valence bond, Z is *tert*-butyl, R<sup>1</sup> = R<sup>2</sup> = R<sup>3</sup> = R<sup>4</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>5</sup> must not be carboxy;

15

when X is -CO-, Y is -O-, A is a valence bond, Z is *tert*-butyl, R<sup>1</sup> = R<sup>2</sup> = R<sup>4</sup> = R<sup>5</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>3</sup> must not be 4-cyanophenyl;

when X is -CO-, the group -Y-A-Z starts with the radical -O-, R<sup>1</sup> = R<sup>2</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>5</sup> must not be carboxy, aminocarbonyl or hydrogen;

20

when -X is -CO-, the group -Y-A-Z starts with the radical -CH<, R<sup>1</sup> = R<sup>2</sup> = R<sup>3</sup> = R<sup>4</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>5</sup> must not be hydroxymethyl, C<sub>1-6</sub>-alkoxy-carbonyl or carboxy; and


25

when X is -CO-, the group -Y-A-Z is 4-methoxyphenyl, R<sup>1</sup> = R<sup>2</sup> = R<sup>3</sup> = R<sup>4</sup> = R<sup>6</sup> = hydrogen, m = n = p = 0 and q = 1, R<sup>5</sup> must not be carboxy;

as well as any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof.

30

In a further aspect the invention relates to novel, substituted imidazoles of the general formula I'''



5

wherein

R<sup>1</sup> is hydrogen or a functional group which can be converted to hydrogen *in vivo*;

10 R<sup>2</sup> is hydrogen, C<sub>1-6</sub>-alkyl, halogen, cyano, trifluoromethyl, hydroxy or -NR<sup>7</sup>R<sup>8</sup>

wherein R<sup>7</sup> and R<sup>8</sup> independently are

hydrogen;

15

C<sub>1-6</sub>-alkyl optionally substituted with aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl amino or C<sub>3-8</sub>-cycloalkyl which are optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl amino;

20

aryl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl amino;

heteroaryl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

- 5 aroyl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

- 10 heteroaroyl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

- 15 arylsulfonyl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

- heteroarylsulfonyl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; or

20

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; or

- 25 R<sup>7</sup> and R<sup>8</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

30

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> independently are



hydrogen; carboxy; C<sub>1-6</sub>-alkoxycarbonyl; -CO-NR<sup>7</sup>R<sup>8</sup> wherein R<sup>7</sup> and R<sup>8</sup> are as defined above; cyano; or halogen;

- 5 C<sub>3-8</sub>-cycloalkyl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

C<sub>1-6</sub>-alkyl optionally substituted with

- 10 C<sub>1-6</sub>-alkoxy; C<sub>1-6</sub>-alkylthio; hydroxy; cyano; halogen; trifluoromethyl; carboxy; C<sub>1-6</sub>-alkoxycarbonyl; or -CO-NR<sup>7</sup>R<sup>8</sup> wherein R<sup>7</sup> and R<sup>8</sup> are as defined above; or C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino which are optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, 15 aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

C<sub>2-6</sub>-alkenyl optionally substituted with

- C<sub>1-6</sub>-alkoxy; C<sub>1-6</sub>-alkylthio; hydroxy; cyano; halogen; trifluoromethyl; carboxy; C<sub>1-6</sub>-alkoxycarbonyl; or -CO-NR<sup>7</sup>R<sup>8</sup> wherein R<sup>7</sup> and R<sup>8</sup> are as defined above; or 20 C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino which are optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

- 25 C<sub>2-6</sub>-alkynyl optionally substituted with

- C<sub>1-6</sub>-alkoxy; C<sub>1-6</sub>-alkylthio; hydroxy; cyano; halogen; trifluoromethyl; carboxy; C<sub>1-6</sub>-alkoxycarbonyl; or -CO-NR<sup>7</sup>R<sup>8</sup> wherein R<sup>7</sup> and R<sup>8</sup> are as defined above; or C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino which are optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, 30 C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; or

aryl optionally substituted with halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, heteroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>7</sup>R<sup>8</sup> wherein R<sup>7</sup> and R<sup>8</sup> are as defined above; or

5

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, and/or R<sup>5</sup> and R<sup>6</sup> together with the carbon atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or hetero-  
 10 arylamino;

m, n, p and q independently are 0, 1 or 2;

15 X is a valence bond, -CH<sub>2</sub>-, -C(=O)-, -C(=S)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)-, -C[=C(CN)<sub>2</sub>]-, -C(=CH-CN)-, or -C(=NR<sup>7</sup>)- wherein R<sup>7</sup> is as defined above;

Y is a valence bond, -O- or -N(R<sup>7</sup>)- wherein R<sup>7</sup> is as defined above;

20

A is a valence bond, C<sub>1-8</sub>-alkylene, C<sub>2-8</sub>-alkenylene, C<sub>2-8</sub>-alkynylene, C<sub>3-8</sub>-cycloalkylene or phenylene; or

when Y is -N(R<sup>7</sup>)-, A may together with R<sup>7</sup> form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and  
 25

Z is -R<sup>9</sup>, -OR<sup>9</sup>, -SR<sup>9</sup>, -NR<sup>9</sup>R<sup>10</sup>, -CHR<sup>9</sup>R<sup>10</sup> or =CR<sup>9</sup>R<sup>10</sup>

30

wherein R<sup>9</sup> and R<sup>10</sup> independently are

hydrogen;

5 C<sub>1-6</sub>-alkyl optionally substituted with aryl, arylamino, heteroaryl, aroyl, heteroaryl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryl-  
oxy, acylamino, heteroaryl or C<sub>3-8</sub>-cycloalkyl which are optionally substituted with  
C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, aryl-  
amino, heteroaryl, aroyl, heteroaryl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkyl-  
sulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino,  
10 halogen, cyano or trifluoromethyl;

C<sub>2-6</sub>-alkenyl optionally substituted with aryl, arylamino, heteroaryl, aroyl, heteroaryl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryl-  
oxy, acylamino, heteroaryl or C<sub>3-8</sub>-cycloalkyl which are optionally substituted with  
15 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, aryl-  
amino, heteroaryl, aroyl, heteroaryl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkyl-  
sulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino,  
halogen, cyano or trifluoromethyl;

20 C<sub>2-6</sub>-alkynyl optionally substituted with aryl, arylamino, heteroaryl, aroyl, heteroaryl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy,  
acylamino, heteroaryl or C<sub>3-8</sub>-cycloalkyl which are optionally substituted with  
C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, aryl-  
amino, heteroaryl, aroyl, heteroaryl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkyl-  
sulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino,  
25 halogen, cyano or trifluoromethyl;

aryl optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl,  
nitro, arylamino, heteroaryl, aroyl, heteroaryl, arylsulfonyl, heteroarylsulfonyl,  
30 C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino,

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl;

- 5 C<sub>3-15</sub>-cycloalkyl optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroaryl-sulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acyl-amino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl;
- 10 aroyl optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl; or
- 15 heteroaryl optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroaryl-sulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl; or
- 20

- R<sup>9</sup> and R<sup>10</sup> are joined by one or more bridging linkers such as a valence bond, C<sub>1-4</sub>-alkylene, C<sub>2-4</sub>-alkenylene, -O-, -S-, -N(R<sup>7</sup>)-, -C(=O)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(R<sup>7</sup>R<sup>8</sup>)-, phenylene, biphenylene, -O-C<sub>1-4</sub>-alkylene, -S-C<sub>1-4</sub>-alkylene, -N(R<sup>7</sup>)-C<sub>1-4</sub>-alkylene,
- 25 -N=C<sub>1-4</sub>-alkylene, -O-C<sub>2-4</sub>-alkenylene, -S-C<sub>2-4</sub>-alkenylene, or -N(R<sup>7</sup>)-C<sub>2-4</sub>-alkenylene, to form a mono-, bi- or polycyclic ring system; or

- when Y is -N(R<sup>7</sup>)-, R<sup>9</sup> or R<sup>10</sup> may together with R<sup>7</sup> form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with
- 30 aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino,

arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl;

with the provisos that

5

when X is -CS-,  $R^1 = R^2 = R^5 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ , the group -Y-A-Z must not start with the radical -NH-;

10

when the group -X-Y-A-Z starts with the radical -CH<sub>2</sub>-,  $R^1 = R^2 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy or aminocarbonyl;

15

when X is -CO-, the group -Y-A-Z starts with the radical -NH-,  $R^1 = R^2 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ , the remainder of the group -Y-A-Z must not be hydrogen, unsubstituted or C<sub>1-6</sub>-alkoxy substituted phenyl, unsubstituted C<sub>3-8</sub>-cycloalkyl or unsubstituted C<sub>1-6</sub>-alkyl;

when X is -CO-, the group -Y-A-Z starts with the radical -O-,  $R^1 = R^2 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy, aminocarbonyl or hydrogen;

20

when -X is -CO-, the group -Y-A-Z starts with the radical -CH<,  $R^1 = R^2 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be hydroxymethyl, C<sub>1-6</sub>-alkoxycarbonyl or carboxy; and

25

when X is -CO-, the group -Y-A-Z is 4-methoxyphenyl,  $R^1 = R^2 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy;

and a pharmaceutically acceptable salt thereof or any optical isomer thereof or mixture of optical isomers, including a racemic mixture, or any tautomeric form.

30

Preferred embodiments thereof are as disclosed above for formula I.

The compounds of the present invention may have one or more asymmetric centres and it is intended that any optical isomers, as separated, pure or partially purified optical isomers or racemic mixtures thereof are included in the scope of the invention.

- 5 Furthermore, geometric isomers may be formed. It is intended that any geometric isomers, as separated, pure or partially purified geometric isomers or mixtures thereof are included within the scope of the invention. Likewise, molecules having a bond with restricted rotation may form geometric isomers. These are also intended to be included within the scope of the present invention.

10

Furthermore, the compounds of the present invention may exist in different tautomeric forms and it is intended that any tautomeric forms which the compounds are able to form are included within the scope of the present invention.

- 15 The present invention also encompasses pharmaceutically acceptable salts of the present compounds. Such salts include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. Acid addition salts include salts of inorganic acids as well as organic acids. Representative examples of suitable inorganic acids include hydrochloric,  
20 hydrobromic, hydroiodic, phosphoric, sulfuric acids and the like. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methanesulfonic, ethanesulfonic, tartaric acids and the like. Further examples of pharmaceutically  
25 acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in *J. Pharm. Sci.* **1977**, *66*, 2, which is incorporated herein by reference. Examples of metal salts include lithium, sodium, potassium, magnesium salts and the like. Examples of ammonium and alkylated ammonium salts include ammonium, methylammonium, dimethylammonium, trimethyl-  
30 ammonium, ethylammonium, hydroxyethylammonium, diethylammonium, butylammonium, tetramethylammonium salts and the like.

Also intended as pharmaceutically acceptable acid addition salts are the hydrates which the present compounds are able to form.

- 5 The acid addition salts may be obtained as the direct products of compound synthesis. In the alternative, the free base may be dissolved in a suitable solvent containing the appropriate acid, and the salt isolated by evaporating the solvent or otherwise separating the salt and solvent.
- 10 The compounds of the present invention may form solvates with standard low molecular weight solvents using methods known to the skilled artisan. Such solvates are also contemplated as being within the scope of the present invention.

The invention also encompasses prodrugs of the present compounds which on administration undergo chemical conversion by metabolic processes before becoming  
15 active pharmacological substances. In general, such prodrugs will be functional derivatives of the compounds which are readily convertible *in vivo* into the present compounds. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H.  
20 Bundgaard, Elsevier, 1985.


The invention also encompasses active metabolites of the present compounds.

The compounds of the present invention interact with the histamine H3 receptor and  
25 may thus be used for the treatment of a wide range of disorders related to the histamine H3 receptor.

Accordingly, in another aspect the present invention relates to a compound of the general formula I as well as any optical or geometric isomer or tautomeric form  
30 thereof including mixtures of these or a pharmaceutically acceptable salt thereof for use as a medicament.

The invention also relates to pharmaceutical compositions comprising, as an active ingredient, at least one compound of the formula I as well as any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof together with one or more pharmaceutically acceptable carriers or diluents.

Furthermore, the invention relates to the use of a compound of the general formula I'



wherein

R<sup>1</sup> is hydrogen or a functional group which can be converted to hydrogen *in vivo*;

R<sup>2</sup> is hydrogen, C<sub>1-6</sub>-alkyl, halogen, cyano, trifluoromethyl, hydroxy or -NR<sup>7</sup>R<sup>8</sup>,

wherein R<sup>7</sup> and R<sup>8</sup> independently are

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with



C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

5 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

10

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

15

R<sup>7</sup> and R<sup>8</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

20

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> independently are

25 hydrogen, carboxy, C<sub>1-6</sub>-alkoxycarbonyl, cyano, trifluoromethyl, halogen,

C<sub>3-8</sub>-cycloalkyl optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

30

C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with  
 C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, cyano, halogen, trifluoromethyl, carboxy,  
 C<sub>1-6</sub>-alkoxycarbonyl,

5 C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or  
 heteroarylamino, which are optionally substituted with  
 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
 trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino,  
 heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

10

aryl optionally substituted with

halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, het-  
 eroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or  
 -CO-NR<sup>9</sup>R<sup>10</sup>,

15

-CO-NR<sup>9</sup>R<sup>10</sup>,

wherein R<sup>9</sup> and R<sup>10</sup> independently are

20

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-  
 amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

25

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
 trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-  
 amino or heteroarylamino,

30

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which  
 are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

5 C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

10 R<sup>9</sup> and R<sup>10</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

15

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, and/or R<sup>5</sup> and R<sup>6</sup> together with the carbon atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

20

25 m, n, p and q independently are 0, 1 or 2;

X is a valence bond, -CH<sub>2</sub>-, -C(=O)-, -C(=S)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)-, -C[=C(CN)<sub>2</sub>]-, -C(=CH-CN)-, -C(=NR<sup>11</sup>)- or -C(=N-S(=O)<sub>2</sub>R<sup>11a</sup>)-,

30 wherein R<sup>11</sup> is

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-  
amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-  
amino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are  
optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino  
or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino  
or heteroarylamino;

R<sup>11a</sup> is

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-  
amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-  
amino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are  
optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

5 Y is a valence bond, -O- or -N(R<sup>12</sup>)-,

wherein R<sup>12</sup> is

hydrogen,

10

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

15

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

20

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

25

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

A is a valence bond, C<sub>1-8</sub>-alkylene, C<sub>2-8</sub>-alkenylene, C<sub>2-8</sub>-alkynylene, C<sub>3-8</sub>-cyclo-alkylene or phenylene, or

30

when Y is  $-N(R^{12})-$ , A, together with  $R^{12}$  and the nitrogen atom to which they are connected, may form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring system optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with  $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxy,  $C_{1-6}$ -alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aryl- $C_{1-6}$ -alkyl, heteroaryl- $C_{1-6}$ -alkyl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and

Z is  $-R^{13}$ ,  $-OR^{13}$ ,  $-SR^{13}$ ,  $-NR^{13}R^{14}$ ,  $-CHR^{13}R^{14}$ ,  $-CR^{13}R^{14}R^{15}$  or  $=CR^{13}R^{14}$ ,

wherein  $R^{13}$ ,  $R^{14}$  and  $R^{15}$  independently are

hydrogen,

$C_{1-6}$ -alkyl,  $C_{2-6}$ -alkenyl or  $C_{2-6}$ -alkynyl, which are optionally substituted with aryl, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl,  $C_{1-6}$ -alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, heteroaryl or  $C_{3-8}$ -cycloalkyl, which are optionally substituted with

$C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxy,  $C_{1-6}$ -alkylthio, aryl- $C_{1-6}$ -alkyl, heteroaryl- $C_{1-6}$ -alkyl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl,  $C_{1-6}$ -alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino, halogen, cyano or trifluoromethyl,

aryl,  $C_{3-15}$ -cycloalkyl, aroyl or heteroaryl, which are optionally substituted with aryl- $C_{1-6}$ -alkyl, heteroaryl- $C_{1-6}$ -alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl,  $C_{1-6}$ -alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino,  $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxy,  $C_{1-6}$ -alkylthio,  $C_{3-8}$ -cycloalkanecarbonyl, hydroxy, amino, halogen, cyano or trifluoromethyl, where

R<sup>13</sup> and R<sup>14</sup> or R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup>, when they do not represent hydrogen, may be joined by one or more bridging linkers such as a valence bond, C<sub>1-4</sub>-alkylene, C<sub>2-4</sub>-alkenylene, -O-, -S-, -N(R<sup>16</sup>)-, -C(=O)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(R<sup>16</sup>R<sup>17</sup>)-, phenylene, biphenylene, -O-C<sub>1-4</sub>-alkylene, -S-C<sub>1-4</sub>-alkylene, -N(R<sup>16</sup>)-C<sub>1-4</sub>-alkylene, -N=C<sub>1-4</sub>-alkylene, -O-C<sub>2-4</sub>-alkenylene, -S-C<sub>2-4</sub>-alkenylene or -N(R<sup>16</sup>)-C<sub>2-4</sub>-alkenylene, to form a mono-, bi- or polycyclic ring system,

wherein R<sup>16</sup> and R<sup>17</sup> independently are

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

R<sup>16</sup> and R<sup>17</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring op-

tionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

as well as any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of disorders related to the histamine H3 receptor.

In still another aspect, the invention relates to a method for the treatment of disorders related to the histamine H3 receptor the method comprising administering to a subject in need thereof an effective amount of a compound of the formula I' as well as any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof or a pharmaceutical composition comprising the same.

More particularly, the present compounds possess histamine H3 receptor antagonistic activity and are accordingly useful in the treatment of a wide range of conditions and disorders in which a histamine H3 receptor blockade is beneficial.

The compounds of the present invention may thus be used for the treatment of airway disorders such as asthma, as anti-diarrhoeals and for the modulation of gastric acid secretion.

The compounds of the present invention may also be used for the treatment of diseases associated with the regulation of sleep and wakefulness and for the treatment of narcolepsy and attention deficit disorders.

Moreover, the compounds of the invention may be used as stimulants or as sedatives.



The compounds of the invention may also be useful for the treatment of eating disorders such as binge eating or bulimia by virtue of their appetite regulating properties.

5

Furthermore, the present compounds may be useful for the treatment and/or prevention of overweight or obesity as well as diseases related to obesity, such as diabetes and cardiovascular disorders.

10 The present compounds may also be used for the treatment of conditions associated with epilepsy. Additionally, the present compounds may be used for the treatment of motion sickness and vertigo, and useful as regulators of hypothalamo-hypophyseal secretion, antidepressants, modulators of cerebral circulation, and in the treatment of irritable bowel syndrome.

15

Further, the compounds of the present invention may be used for the treatment of dementia and Alzheimer's disease. Moreover, the compounds of the present invention may be used as analgetics and for the treatment of inflammatory painful conditions or neurogenic inflammation.

20

These new compounds may also interact with the vanilloid receptors, the serotonin receptors, and the adrenergic receptors and may be useful for the treatment of diseases associated with these receptors. Hence, the compounds of the present invention may be vanilloid receptor agonists, and thus be useful for the treatment of  
25 obesity by enhancement of the metabolic rate and energy expenditure. Further, by virtue of their interaction with the vanilloid receptor the compounds of the present invention may be useful for the treatment of pain or neurogenic inflammation or inflammatory painful conditions.

30 Furthermore, by virtue of their interaction with the 5-HT<sub>3</sub> receptor (serotonin-3-receptor), the compounds of the present invention may be useful as antiemetics, in

particular the chemotherapy-induced emesis. Further potential applications of 5-HT<sub>3</sub> antagonists include treatment of central nervous system disorders such as anxiety, schizophrenia, drug abuse and withdrawal symptoms, and pathological and age-associated amnesia.

5

Furthermore, the present compounds may be alpha-2-adrenoceptor agonists or antagonists and thus be useful for the treatment of hypertension and of conditions associated with overexpression or hypersensitization of adrenergic alpha-2 receptors, especially obesity, withdrawal symptoms to an adrenergic alpha-2 agonist, neurological disorders (especially orthostatic hypotension), multiple system atrophy, diabetes mellitus, benign prostatic hyperplasia or drug induced sensitization of adrenergic alpha-2 receptors. Moreover, the compounds of the present invention, by virtue of their interaction with alpha-2 receptors, may be useful as sedatives and hypnotics (sleep inducing agents) or as stimulants.

15

In a preferred embodiment of the invention the present compounds are used for the preparation of a medicament for the treatment and/or prevention of overweight or obesity.

20 In another preferred embodiment of the invention the present compounds are used for the preparation of a medicament for the suppression of appetite or satiety induction.

In a further preferred embodiment of the invention the present compounds are used for the preparation of a medicament for the prevention and/or treatment of disorders and diseases related to overweight or obesity such as as atherosclerosis, hypertension, diabetes, especially Type 2 diabetes (NIDDM (non-insulin dependent diabetes mellitus)), dyslipidaemia, coronary heart disease, gallbladder disease, osteoarthritis and various types of cancer such as endometrial, breast, prostate and colon cancers.

30

In yet a further preferred embodiment of the invention the present compounds are used for the preparation of a medicament for the prevention and/or treatment of eating disorders such as bulimia and binge eating.

- 5 In a further aspect of the invention the present compounds may be administered in combination with one or more further pharmacologically active substances eg other antiobesity agents or appetite regulating agents.

Such agents may be selected from the group consisting of CART agonists, NPY  
10 antagonists, MC4 agonists, orexin antagonists, TNF agonists, CRF agonists, CRF BP antagonists, urocortin agonists,  $\beta$ 3 agonists, MSH (melanocyte-stimulating hormone) agonists, MCH (melanocyte-concentrating hormone) antagonists, CCK agonists, serotonin re-uptake inhibitors, mixed serotonin and noradrenergic compounds, 5HT agonists, bombesin agonists, galanin antagonists, growth hormone, growth hormone  
15 releasing compounds, TRH agonists, uncoupling protein 2 or 3 modulators, GLP-1, leptin agonists, DA agonists (bromocriptin, doprexin), lipase/amylase inhibitors, PPAR modulators, RXR modulators or TR  $\beta$  agonists.

In a preferred embodiment of the invention the antiobesity agent is leptin.

20

In another preferred embodiment the antiobesity agent is dexamphetamine or amphetamine.

In another preferred embodiment the antiobesity agent is fenfluramine or  
25 dexfenfluramine.

In still another preferred embodiment the antiobesity agent is sibutramine.

In a further preferred embodiment the antiobesity agent is orlistat.

30

In another preferred embodiment the antiobesity agent is mazindol or phentermine.

The present compounds may also be administered in combination with one or more antidiabetics or other pharmacologically active material(s), including compounds for the treatment and/or prophylaxis of insulin resistance and diseases, wherein insulin resistance is the pathophysiological mechanism. Suitable antidiabetics comprise insulin, GLP-1 derivatives such as those disclosed in WO 98/08871 to Novo Nordisk A/S which is incorporated herein by reference as well as orally active hypoglycaemic agents.

10 The orally active hypoglycaemic agents preferably comprise sulphonylureas, biguanides, meglitinides, oxadiazolidinediones, thiazolidinediones, glucosidase inhibitors, glucagon antagonists such as those disclosed in WO 99/01423 to Novo Nordisk A/S and Agouron Pharmaceuticals, Inc., GLP-1 agonists, potassium channel openers such as those disclosed in WO 97/26265 and WO 99/03861 to  
15 Novo Nordisk A/S which are incorporated herein by reference, insulin sensitizers such as those disclosed in WO 99/19313 to Dr. Reddy's Research Foundation, DPP-IV inhibitors, inhibitors of hepatic enzymes involved in stimulation of gluconeogenesis and/or glycogenolysis, glucose uptake modulators, compounds modifying the lipid metabolism such as antihyperlipidemic agents and antilipidemic agents, compounds  
20 lowering food intake, PPAR and RXR agonists and agents acting on the ATP-dependent potassium channel of the  $\beta$ -cells.

In a preferred embodiment of the invention the present compounds are administered in combination with insulin.

25

In a further preferred embodiment the present compounds are administered in combination with a sulphonylurea eg tolbutamide, glibenclamide, glipizide or glicazide.

30 In another preferred embodiment the present compounds are administered in combination with a biguanide eg metformin.

In yet another embodiment the present compounds are administered in combination with a meglitinide eg repaglinide.

- 5 In still another preferred embodiment the present compounds are administered in combination with a thiazolidinedione selected from troglitazone, ciglitazone, pioglitazone, rosiglitazone and the compounds disclosed in WO 97/41097 to Dr. Reddy's Research Foundation, especially 5-[[4-[(3,4-dihydro-3-methyl-4-oxo-2-quinazolinylmethoxy]phenyl]-methyl]-2,4-thiazolidinedione.

10

In a further preferred embodiment the present compounds are administered in combination with an  $\alpha$ -glucosidase inhibitor eg miglitol or acarbose.

- 15 In yet a preferred embodiment the present compounds are administered in combination with an agent acting on the ATP-dependent potassium channel of the  $\beta$ -cells selected from tolbutamide, glibenclamide, glipizide, glicazide and repaglinide.

Furthermore, the present compounds may be administered in combination with nateglinide.

20

In still another embodiment the present compounds are administered in combination with an antihyperlipidemic agent or antilipidemic agent eg cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

25

- 30 In a further embodiment the present compounds are administered in combination with more than one of the above-mentioned compounds eg in combination with a sulphonylurea and metformin, a sulphonylurea and acarbose, repaglinide and metformin, insulin and a sulphonylurea, insulin and metformin, insulin and troglitazone, insulin and lovastatin, etc.

Furthermore, the present compounds may be administered in combination with one or more antihypertensive agent. Examples of antihypertensive agents are  $\beta$ -blockers such as alprenolol, atenolol, timolol, pindolol, propranolol and metoprolol, ACE  
5 (angiotensin converting enzyme) inhibitors such as benazepril, captopril, enalapril, fosinopril, lisinopril, quinapril and ramipril, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, nimodipine, diltiazem and verapamil, and  $\alpha$ -blockers such as doxazosin, urapidil, prazosin and terazosin. Further  
reference can be made to Remington: The Science and Practice of Pharmacy, 19<sup>th</sup>  
10 Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

It should be understood that any suitable combination of the compounds according to the invention with one or more of the above-mentioned compounds and optionally one or more further pharmacologically active substances are considered to be within  
15 the scope of the present invention.

### PHARMACEUTICAL COMPOSITIONS

The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or excipients, in either single or multiple doses.  
20 The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19<sup>th</sup> Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

25 The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal)  
30 ermal) route, the oral route being preferred. It will be appreciated that the preferred

route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient chosen.

Pharmaceutical compositions for oral administration include solid dosage forms such as capsules, tablets, dragees, pills, lozenges, powders and granules. Where appropriate, they can be prepared with coatings such as enteric coatings or they can be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well-known in the art.

Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

Pharmaceutical compositions for parenteral administration include sterile aqueous and non-aqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

Other suitable administration forms include suppositories, sprays, ointments, cremes, gels, inhalants, dermal patches, implants etc.

A typical oral dosage is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. A typical unit dosage form for oral administration one or more times per day such as 1 to 3 times per day may contain of from 0.05 to about 1000 mg, preferably from about 0.1 to about 500 mg, and more preferred from about 0.5 mg to about 200 mg.

For parenteral routes, such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for oral administration.

10

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. One example is an acid addition salt of a compound having the utility of a free base. When a compound according to the invention contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of a free base of the compound according to the invention with a chemical equivalent of a pharmaceutically acceptable acid, for example, inorganic and organic acids. Representative examples are mentioned above. Physiologically acceptable salts of a compound with a hydroxy group include the anion of said compound in combination with a suitable cation such as sodium or ammonium ion.

20

For parenteral administration, solutions of the present compounds in sterile aqueous solution, aqueous propylene glycol or sesame or peanut oil may be employed. Such aqueous solutions should be suitable buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

25

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatine, agar, pectin, acacia, magnesium stearate,

30



stearic acid or lower alkyl ethers of cellulose. Examples of liquid carriers are syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene or water. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or  
 5 mixed with a wax. The pharmaceutical compositions formed by combining the compounds according to the invention and the pharmaceutically acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

10 Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and which may include a suitable excipient. These formulations may be in the form of powder or granules, as a solution or suspension in an  
 15 aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion.

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatine capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from  
 20 about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

A typical tablet which may be prepared by conventional tableting techniques may  
 25 contain:

Core:

Active compound (as free compound or salt thereof)	5.0 mg
Lactosum Ph. Eur.	67.8 mg
30 Cellulose, microcryst. (Avicel)	31.4 mg
Amberlite	1.0 mg

Magnesii stearas Ph. Eur.

q.s.

Coating:

HPMC approx.

9 mg

5 Mywacett 9-40 T\* approx.


0.9 mg

\*Acylated monoglyceride used as plasticizer for film coating.


10 If desired, the pharmaceutical composition of the invention may comprise the compound of the formula I' in combination with further pharmacologically active substances.

The preparation of the compounds of this invention can be realized in many different ways. The preparation of imidazoles of the formula III has been described in the literature (see e.g. F. B. Stocker et al., *J. Org. Chem.* **1966**, 31, 2380; idem, *ibid.* **1990**, 55, 3370; T. Vitali et al., *Il Farmaco* **1967**, 22, 821; idem, *ibid.* **1965**, 20, 634; S. Fränkel, K. Zeimer, *Biochemische Zeitschrift* **1920**, 110, 238; G. Arcari et al., *Fr. Pat.* 1976, 2 337 726; DE 2700012, 1977, *Chem. Abstr.*, 87, 201535).

20 Compounds of the formula Ia, wherein R<sup>1</sup> and R<sup>2</sup> are hydrogen, m = n = p = 0 and q = 1 can be prepared as outlined below:



25 More specifically, different types of compounds of the formula Ia of this invention can be prepared by the methods 1) to 4) sketched below:



LG = leaving group

For instance, compounds of the formula **Ib** can be synthesized from 4,5,6,7-tetrahydroimidazo[4,5-c]pyridines **III** by treating the latter with suitable activated derivatives of carboxylic acids, such as acyl imidazoles, anhydrides, acid chlorides or active esters, or any of the derivatives commonly used for the preparation of carboxamides, under appropriate conditions.

Compounds of the formula **Ic** can be prepared by treating 4,5,6,7-tetrahydroimidazo[4,5-c]pyridines **III** with isocyanates Z-A-NCO or with synthetic equivalents thereof, such as carbamoyl chlorides Z-A-N(R<sup>12</sup>)-CO-Cl under suitable conditions.

Compounds of the formula **Id** can be synthesized by treating 4,5,6,7-tetrahydro-imidazo[4,5-c]pyridines **III** with haloformates Z-A-O-CO-Cl or with synthetic equivalents thereof such as activated carbonates (e.g. 4-nitrophenyl carbonates) under suitable conditions.

Finally, compounds of the formula **Ie** can be prepared by treating 4,5,6,7-tetrahydro-imidazo[4,5-c]pyridines **III** with isothiocyanates Z-A-NCS or with synthetic equivalents thereof, such as thiocarbamoyl chlorides Z-A-N(R<sup>12</sup>)-CS-Cl under suitable conditions.

The starting materials are either known compounds or compounds which may be prepared in analogy with the preparation of similar known compounds.

The present invention is further illustrated by the following representative examples, which are, however, not intended to limit the scope of the invention in any way.

## EXAMPLES

In the examples the following terms are intended to have the following, general meanings:

DCM: dichloromethane, methylenechloride

DMF: N,N-dimethyl formamide

DMSO: dimethyl sulfoxide

EDC: N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride

HOBt: N-hydroxybenzotriazole, 1-hydroxybenzotriazole

NMP: N-methylpyrrolidone

NMR spectra were recorded on Bruker 300 MHz and 400 MHz instruments. HPLC-MS was performed on a Perkin Elmer instrument (API 100).


HPLC-systems from Merck-Hitachi (Hibar™ RT 250-4, Lichrosorb™ RP 18, 5.0 μm, 4.0 x 250 mm, gradient elution, 20% to 80% acetonitrile in water within 30 min, 1.0 mL/min, detection at 254 nm) and Waters (Symmetry™, C<sub>18</sub>, 3.5 μm, 3.0 x 150 mm, gradient elution, 5% to 90% acetonitrile in water within 15 min, 1.0 mL/min, detection at 214 nm) were used.

Furthermore, where stated the following HPLC method h8 was used:

The reverse phase analysis was performed using UV detections at 214, 254, 276 and 301 nm on a 218TP54 4.6 mm x 150 mm C-18 silica column, which was eluted at 1 mL/min at 42 °C. The column was equilibrated with 5% acetonitrile, 85% water and 10% of a solution of 0.5% trifluoroacetic acid in water and eluted by a linear gradient from 5% acetonitrile, 85% water and 10% of a solution of 0.5% trifluoroacetic acid to 90% acetonitrile and 10% of a solution of 0.5% trifluoroacetic acid over 15 min.

Where stated the following general procedures were used:

#### General Procedure A:



20

To a solution of the dihydrochloride of the amine II (10.0 mmol) in water (5 mL) an aqueous sodium hydroxide solution (12 N, 4.2 mL, 50.4 mmol), methanol (35 mL) and the carbonyl compound R<sup>3</sup>R<sup>4</sup>CO (25.0 mmol) were added. The resulting mixture was refluxed overnight, water (15 mL) was added, methanol was evaporated under reduced pressure and the residue was diluted with water to a volume of approximately 25 mL. The resulting mixture was washed with ether (2 x 50 mL, removal of excess ketone) and then, while stirring vigorously, the acyl halide (Z-A-Y-CO-Cl, 11.5

25


mmol) was added portionwise. After stirring for 10 min the mixture was extracted with DCM (2 x 100 mL). The combined organic phases were washed with water (25 mL), dried ( $\text{MgSO}_4$ ) and concentrated. The remaining oil was redissolved in ethyl acetate (100 mL) and a solution of oxalic acid (0.45 g, 5.0 mmol) in ethyl acetate (25 mL) was added. After stirring for 30 min the precipitate was filtered off and dried under reduced pressure. Alternatively, the crude product **1b** could be purified by column chromatography (silica gel, gradient elution with heptane/ethyl acetate/methanol).

#### General Procedure B:

A mixture of the dihydrochloride of the amine **II** (95 mmol), water (200 mL), and the carbonyl compound  $\text{R}^3\text{R}^4\text{CO}$  (133 mmol) was refluxed until no more amine **II** could be detected (HPLC). The mixture was then concentrated to dryness and the crude product **III** was purified by recrystallization.

The purified amine **III** was then acylated as in General Procedure A or by any other, conventional method.

**Example 1:** 5-(3-Cyclohexylpropanoyl)-4,4-dimethyl-4,5,6,7-tetrahydroimidazo-[4,5-c]pyridine oxalic acid salt




Following the General Procedure A using histamine dihydrochloride (1.84 g, 10.0 mmol), acetone (4.0 mL) and 3-cyclohexylpropanoyl chloride (2.0 g, 11.5 mmol) 0.80 g (21%) of the title amide was obtained as oxalic acid salt.

HPLC (214 nm): elution at 18.59 min. LC-MS: Calcd. for  $\text{MH}^+$ : 290; found: 290.

5390.029-DK,

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>, two rotamers, 6:4): δ 0.80-0.95 (m, 2H), 1.05-1.28 (m, 4H), 1.35-1.76 (m, 13H), 2.38 (m, 1.2H), 2.64 (t, *J* = 5 Hz, 1.2H), 2.89 (t, *J* = 5 Hz, 0.8H), 3.01 (t, *J* = 7 Hz, 0.8H), 3.54 (t, *J* = 5 Hz, 0.8H), 3.58 (t, *J* = 5 Hz, 1.2H), 8.25 (s, 0.6H), 8.34 (s, 0.4H).

**Example 2:** 5-(5-Cyclohexylpentanoyl)-4,5,6,7-tetrahydroimidazo[4,5-*c*]pyridine




10

To a solution of 5-cyclohexylpentanoic acid (0.94 g, 5.10 mmol) in DCM (4 mL) carbonyldiimidazole (0.83 g, 5.12 mmol) was added. The resulting mixture was stirred at room temperature for 16 h and then added to a solution of 4,5,6,7-tetrahydroimidazo[4,5-*c*]pyridine (5.10 mmol) in DCM (4 mL). After 2.5 h DCM (50 mL) was added and the mixture was washed with water (3 x 15 mL). The organic layer was then dried (MgSO<sub>4</sub>) and concentrated. The crude product was purified by column chromatography (silica gel, ethyl acetate/methanol 9:1) whereby 0.35 g (24%) of the title amide was obtained as an oil.

20 HPLC (214 nm): elution at 20.74 min. LC-MS: Calcd. for MH<sup>+</sup>: 290; found: 290.


<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>, two rotamers, 1:1): δ 0.70-0.80 (m, 2H), 1.05-1.34 (m, 8H), 1.43-1.52 (m, 2H), 1.55-1.71 (m, 5H), 2.36 (m, 2H), 2.52 (t, *J* = 5 Hz, 1H), 2.62 (t, *J* = 5 Hz, 1H), 3.68 (t, *J* = 5 Hz, 1H), 3.74 (t, *J* = 5 Hz, 1H), 4.41 (s, 2H), 7.47 (s, 0.5H), 7.49 (s, 0.5H), 11.85 (s, br, 1H).

25

**Example 3: 5-(6-Phenylhexanoyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine**

- 5 To a solution of carbonyldiimidazole (0.83 g, 5.12 mmol) in DCM (8 mL) 6-phenylhexanoic acid (0.98 g, 5.10 mmol) was added dropwise. The mixture was stirred at room temperature for 20 h, and then a solution of 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine (5.10 mmol) in DMF (1 mL) and DCM (1 mL) was added. After stirring for four days DCM (100 mL) was added and the mixture was washed with water (15 mL) and dried (MgSO<sub>4</sub>). Concentration and column chromatography (silica gel, ethyl acetate/methanol 9:1) gave 0.80 g (53%) of the title amide as an oil.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>, two rotamers, 1:1):  $\delta$  1.22-1.38 (m, 2H), 1.46-1.64 (m, 4H), 2.38 (m, 2H), 2.45-2.68 (m, 6H), 3.68 (t, *J* = 5 Hz, 1H), 3.72 (t, *J* = 5 Hz, 1H), 4.41 (s, 2H), 7.14-7.29 (m, 5H), 7.48 (s, 0.5H), 7.50 (s, 0.5H), 11.90 (s, br, 1H).

**Example 4: 5-(3-Cyclohexylpropanoyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine**

20

- To a solution of carbonyldiimidazole (0.83 g, 5.12 mmol) in DCM (8 mL) 3-cyclohexylpropionic acid (0.80 g, 5.10 mmol) was added dropwise. The mixture was stirred at room temperature for 20 h, and then a solution of 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine (5.10 mmol) in DMF (1 mL) and DCM (1 mL) was
- 25 5390.029-DK,




added. After stirring for four days DCM (100 mL) was added and the mixture was washed with water (15 mL) and dried (MgSO<sub>4</sub>). Concentration and column chromatography (silica gel, ethyl acetate/methanol 9:1) gave 0.66 g (50%) of the title amide as an oil.

5

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>, two rotamers, 1:1):  $\delta$  0.80-0.95 (m, 2H), 1.05-1.30 (m, 4H), 1.90 (m, 2H), 1.56-1.75 (m, 5H), 2.39 (m, 2H), 2.52 (t, *J* = 5 Hz, 1H), 2.61 (t, *J* = 5 Hz, 1H), 3.69 (t, *J* = 5 Hz, 1H), 3.72 (t, *J* = 5 Hz, 1H), 4.42 (s, 2H), 7.51 (s, 0.5H), 7.53 (s, 0.5H), 11.90 (s, br, 1H).

10

**Example 5:** 5-[3-(4-Fluorophenyl)propanoyl]-4-(4-isopropylphenyl)-4,5,6,7-tetrahydroimidazo[4,5-*c*]pyridine



15

A mixture of histamine dihydrochloride (1.85 g, 10.0 mmol), water (10 mL), potassium hydroxide (1.72 g, 30.0 mmol), ethanol (25 mL) and 4-isopropylbenzaldehyde (1.62 g, 10.91 mmol) was heated to reflux for 1.5 h. Ethanol was evaporated and the residue was diluted with water (40 mL). Extraction (5 x 25 mL DCM), washing of the combined extracts (2 x 50 mL brine) and drying (MgSO<sub>4</sub>) yielded 2.29 g (87%) of crude 4-(4-isopropylphenyl)-4,5,6,7-tetrahydro-imidazo[4,5-*c*]pyridine, which was used for the next synthetic step without further purification. This amine (0.48 g, 1.99 mmol) was dissolved in DCM (5 mL) and added to a 30 min old mixture of 3-(4-fluorophenyl)propionic acid (0.31 g, 1.84 mmol), HOBt (0.27 g, 1.20 mmol) and EDC (0.42 g, 2.19 mmol) in DCM (10 mL). After 18 h the mixture was washed with water (50 mL), dried (MgSO<sub>4</sub>) and concentrated. The crude product was purified by column 5390.029-DK,

20

25


chromatography (silica gel, gradient elution with DCM/methanol). 0.24 g (33%) of the title amide was obtained.

HPLC (214 nm): elution at 10.21 min. LC-MS: Calcd. for  $MH^+$ : 392; found: 392.

5

$^1H$  NMR (400 MHz,  $DMSO-d_6$ ):  $\delta$  1.18 (d,  $J = 7$  Hz, 6H), 2.50-2.94 (m, 7H), 3.05 (m, 1H), 3.95 (m, 1H), 6.48 (s, br, 0.7H), 6.67 (s, br, 0.3H), 6.99-7.35 (m, 8H), 7.55 (s, 1H), 11.90 (s, 1H).

10 **Example 6:** 5-(3,3-Diphenylpropanoyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine




To a suspension of 3,3-diphenylpropionic acid (14 mg, 0.06 mmol) and HOBt (9 mg, 0.07 mmol) in ethyl acetate (1.5 mL) a solution of EDC (12 mg, 0.06 mmol) in ethyl acetate (0.5 mL) was added. The resulting mixture was shaken for 20 min at room temperature and then 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine dihydrochloride (12 mg, 0.06 mmol) and triethylamine (0.02 mL) were added. After shaking for 16 h the mixture was washed with brine (2 x 2 mL), and the organic phase was concentrated.  
12 mg (60%) of the title amide was obtained.

HPLC (214 nm): elution at 8.71 min. LC-MS: Calcd. for  $MH^+$ : 332; found: 332.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ , two rotamers, 1:1):  $\delta$  2.55 (m, 2H), 3.15 (t,  $J = 7$  Hz, 2H), 3.61 (t,  $J = 5$  Hz, 1H), 3.80 (t,  $J = 5$  Hz, 1H), 4.41 (s, 1H), 4.57 (s, 1H), 4.67 (m, 1H), 7.06-7.30 (m, 10H), 7.39 (s, 0.5H), 7.43 (s, 0.5H).

- 5 **Example 7:** 5-(3-Cyclohexylpropanoyl)-4-ethyl-4-methyl-4,5,6,7-tetrahydroimidazo-[4,5-c]pyridine oxalic acid salt



- 10 From histamine dihydrochloride (1.85 g, 10.0 mmol), 2-butanone (1.80 g, 25.0 mmol) and 3-cyclohexylpropanoyl chloride (2.0 g, 11.5 mmol) 0.60 g (15%) of the title oxalate was obtained using the General Procedure A.


HPLC (214 nm): elution at 19.54 min. LC-MS: Calcd. for  $\text{MH}^+$ : 304; found: 304.

15

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ , two rotamers, 2:1):  $\delta$  0.39 (t,  $J = 7$  Hz, 2H), 0.79-0.93 (m, 2H), 0.95 (t,  $J = 7$  Hz, 1H), 1.05-1.29 (m, 4H), 1.39 (m, 2H), 1.49 (s, 1H), 1.59 (s, 2H), 1.60-2.10 (m, 7H), 2.15-2.50 (m, 1H), 2.60-2.99 (m, 3H), 3.28-3.49 (m, 1.3H), 3.88 (m, 0.7H), 7.69 (s, 0.3H), 8.39 (s, 0.7H).

20

**Example 8:** 5-[3-(10,11-Dihydro-5*H*-dibenzo[*b,f*]azepin-5-yl)propanoyl]-4,5,6,7-tetrahydroimidazo[4,5-*c*]pyridine



5

Iminodibenzyl (50.0 g, 0.256 mol) was dissolved in DMF (700 mL), sodium hydride (12.3 g, 0.306 mol, 60% dispersion in oil) was slowly added in portions and the mixture was stirred at 50 °C for 2 h. Ethyl 3-bromopropionate (100 mL, 0.77 mol) was slowly added dropwise and the mixture was heated at reflux temperature overnight.

10 The mixture was cooled and evaporated. The residue was suspended in DCM (150 mL), filtered and the solvent was evaporated. The resulting residue was purified in portions by column chromatography (silica gel, DCM) to give 5.1 g (7%) of 3-(10,11-dihydro-5*H*-dibenzo[*b,f*]azepin-5-yl)propionic acid ethyl ester.

15 TLC:  $R_f$  = 0.69 (silica gel, DCM).

3-(10,11-Dihydro-5*H*-dibenzo[*b,f*]azepin-5-yl)propionic acid ethyl ester (1.41 g, 4.77 mmol) was dissolved in ethanol (30 mL) and a solution of sodium hydroxide (0.75 g, 18.8 mmol) in water (5 mL) was added. The mixture was stirred for 3.5 h. 1 N Hydrochloric acid (17 mL) was added and the mixture was extracted with DCM (2 x 25 mL).  
20 The combined organic extracts were washed with brine (50 mL), dried ( $\text{MgSO}_4$ ) and the solvent was evaporated to give 1.18 g (92%) of 3-(10,11-dihydro-5*H*-dibenzo[*b,f*]azepin-5-yl)propionic acid.


25 Carbonyldiimidazole (0.33 g, 2.1 mmol) was dissolved in DCM (5 mL) and 3-(10,11-dihydro-5*H*-dibenzo[*b,f*]azepin-5-yl)-1-propionic acid (0.56 g, 2.1 mmol) was added.

The mixture was stirred at room temperature for 1.5 h under a nitrogen atmosphere. Simultaneously, sodium methoxide (0.8 mL of a 30% solution in water, 4.4 mmol) was added to a solution of 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine dihydrochloride (0.43 g, 2.2 mmol) in methanol (5 mL). The mixture was stirred at room temperature  
 5 for 1.5 h under a nitrogen atmosphere. The solvent was evaporated and the residue was stripped with DCM (6 mL). The above solution of the activated carboxylic acid was added to the residue and the reaction mixture was stirred at room temperature overnight. Water (10 mL) was added followed by DCM (50 mL) and the phases were separated. The aqueous phase was extracted with DCM (20 mL) and the combined  
 10 organic phases were dried ( $\text{MgSO}_4$ ). After evaporation of the solvent, the residue was purified by column chromatography (silica gel, 150 mL, methanol/ ethyl acetate 1:5). Evaporation of the solvent afforded 0.41 g (52%) of the title compound as a solid.

15 TLC:  $R_f$  = 0.28 (silica gel, methanol/ethyl acetate 1:5). LC-MS: Calcd. for  $\text{MH}^+$ : 373; found: 373.

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ , two rotamers, 1:1):  $\delta$  2.49 (m, 1H), 2.60-2.72 (m, 3H), 3.07 (d, 4H), 3.47 (t, 1H), 3.86 (t, 1H), 4.08-4.20 (m, 2H), 4.22 (s, 1H), 4.62 (s, 1H),  
 20 6.90 (m, 2H), 7.01-7.16 (m, 6H), 7.40 (s, 0.5H), 7.45 (s, 0.5H).

**Example 9:** 5-(3-Cyclohexylpropanoyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine-4-spirocyclobutane oxalic acid salt




From histamine dihydrochloride (1.85 g, 10.0 mmol), cyclobutanone (1.75 g, 25.0 mmol) and 3-cyclohexylpropanoyl chloride (2.0 g, 11.5 mmol), 0.70 g (18%) of the title oxalate was obtained (General Procedure A).

5 HPLC (214 nm): elution at 19.40 min. LC-MS: Calcd. for  $MH^+$ : 302; found: 302.

$^1H$  NMR (400 MHz,  $DMSO-d_6$ ):  $\delta$  0.75-0.95 (m, 2H), 1.04-1.28 (m, 4H), 1.35 (q,  $J$  = 7 Hz, 2H), 1.50-2.70 (m, 11H), 3.63 (t,  $J$  = 5 Hz, 2H), 8.22 (s, br, 1H).

### 10 Example 10: Parallel Synthesis of Ureas

To each reactor of an array of 12 reactors, each containing a solution of 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine dihydrochloride (0.07 mmol) in DMF (0.5 mL, containing 5% triethylamine) a solution of one isocyanate (0.9 equivalents) selected from 12 different isocyanates in 1,2-dichloroethane (0.2 mL) was added. The resulting  
 15 mixtures were shaken overnight at room temperature. Concentration under reduced pressure gave the corresponding ureas. Using this methodology the following ureas were prepared:




20

Example	Z-A-	$MH^+$ (calcd)	$MH^+$ (found)
10-001	2-(2-thienyl)ethyl	277	277
10-002	3,5-dimethyl-1,2-oxazol-4-yl	262	262
10-003	1-(1-naphthyl)ethyl	321	
10-004	(2-phenylcyclopropyl)	283	283
10-005	1-(4-bromophenyl)ethyl	350	
10-006	2-(trifluoromethyl)phenyl	311	311

10-007	2-phenylethyl	271	271
10-008	4-(trifluoromethyl)phenyl	311	311
10-009	3-cyanophenyl	268	268
10-010	4-cyanophenyl	268	268
10-011	n-octyl	279	279

**Example 11:** 5-(2,4-Dichlorobenzylaminocarbonyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine



5

To a mixture of 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine dihydrochloride (0.50 g, 2.55 mmol), ethanol (10 mL) and triethylamine (1.10 mL, 7.89 mmol), 2,4-dichlorobenzyl isocyanate (0.52 g, 2.57 mmol) was added. The resulting mixture was stirred at room temperature for 16 h and then concentrated under reduced pressure. The residue was distributed between water (20 mL) and ethyl acetate (75 mL). The organic phase was washed with water (10 mL) and brine (10 mL), dried (MgSO<sub>4</sub>) and concentrated under reduced pressure. The residue was triturated in diethylether (50 mL) and the crude product was filtered off. Recrystallization from acetone (25 mL) gave 0.30 g (37%) of the title urea as a colourless solid.


15

HPLC (214 nm): elution at 7.97 min. LC-MS: Calcd. for MH<sup>+</sup>: 325; found: 325.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.56 (s, br, 2H), 3.65 (t, *J* = 6 Hz, 2H), 4.29 (d, *J* = 6 Hz, 2H), 4.34 (s, 2H), 7.20 (s, br, 1H), 7.29 (d, *J* = 8 Hz, 1H), 7.39 (dd, *J* = 8, 1 Hz, 1H), 7.47 (s, 1H), 7.56 (d, *J* = 1 Hz, 1H), 11.79 (s, br, 1H).

20

**Example 12:** 4-(2-Cyclohexylethyl)-5-(ethylaminocarbonyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine



5

To a solution of histamine dihydrochloride (1.84 g, 10.0 mmol) in water (5 mL) methanol (40 mL) was added. To this stirred mixture an aqueous 12 N sodium hydroxide solution (4.2 mL, 50.4 mmol) and 3-(cyclohexyl)propionaldehyde (3.5 g, 25 mmol) were added. The resulting mixture was heated at reflux temperature for 20 h.

10 Concentrated hydrochloric acid was added until pH 1 and the mixture was diluted with water (100 mL). The mixture was washed with diethyl ether (3 x 25 mL) and the aqueous solution was concentrated under reduced pressure. The residue was suspended in methanol (150 mL) and the suspension was filtered. The filtrate was concentrated under reduced pressure to give a residue which was treated with warm

15 ethanol (75 mL). The resulting mixture was filtered and the filtrate was concentrated under reduced pressure to a volume of approximately 10 mL. Acetone (50 mL) was added and the mixture was left for crystallization. The solid was isolated by filtration and dried under reduced pressure to give 2.4 g of a solid which was dissolved in water (10 mL). While stirring, an aqueous 1 N sodium hydroxide solution was added


20 until pH 11-12. The resulting mixture was extracted with ethyl acetate (200 mL), the extracts were dried ( $\text{MgSO}_4$ ) and the solvent was evaporated under reduced pressure. This afforded 1.25 g of 4-(2-cyclohexylethyl)-4,5,6,7-tetrahydroimidazo[4,5-c]-pyridine.



To a solution of the above amine (0.45 g, 1.9 mmol) in ethanol (10 mL), triethylamine (0.2 g, 1.9 mmol) and ethyl isocyanate (0.14 g, 1.9 mmol) were added. The mixture was stirred overnight at room temperature and then concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, elution with ethyl acetate/methanol 9:1). This afforded 0.14 g (24% calculated from the amine) of the title compound as a solid.

M.p. 210-212 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 0.85 (m, 2H), 1.00 (t, 3H), 1.05-1.45 (m, 6H), 1.50-1.75 (m, 7H), 2.35 (m, 1H), 2.55 (m, 1H), 3.03 (m, 3H), 4.10 (m, br, 1H), 4.85 (m, br, 1H), 6.38 (s, br, 1H), 7.40 (s, 1H), 11.68 (s, br, 1H).


**Example 13:** 4-(2-Cyclohexylethyl)-5-(2,4-dichlorobenzylaminocarbonyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine









To a solution of 4-(2-cyclohexylethyl)-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine (0.47 g, 2.0 mmol, prepared as described in Example 13) in ethanol (10 mL), triethylamine (0.28 mL, 2.0 mmol) and 2,4-dichlorobenzyl isocyanate (0.41 g, 2.0 mmol) were added dropwise. The mixture was stirred overnight at room temperature and then concentrated under reduced pressure. The residue was re-evaporated with acetone and then dissolved in acetone (15 mL) and left for crystallization. The solid was isolated by filtration, washed with acetone and dried. This afforded 0.60 g (69%) of the title compound as a solid.

M.p. 185-187 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 0.85 (m, 2H), 1.05-1.45 (m, 6H), 1.52-1.70 (m, 7H), 2.42 (m, 1H), 2.60 (m, 1H), 3.09 (m, 1H), 4.18 (m, br, 1H), 4.28 (m, 2H), 4.96 (m, br, 1H), 7.10 (t, br, 1H), 7.28 (d, 1H), 7.38 (dd, 1H), 7.43 (s, 1H), 7.55 (d, 1H), 11.75 (s, br, 1H).

#### Example 14: Parallel Synthesis of Carboxamides




To each reactor in an array of six, a suspension of HOBt (7.4 mg, 55 μmol) in a mixture of acetonitrile, 1,2-dichloroethane, NMP and DMSO (250 μL) was added. Then a suspension of EDC (11.5 mg, 60 μmol) in a mixture of acetonitrile, 1,2-dichloroethane, NMP and DMSO (250 μL) was added to each reactor. To each reactor a carboxylic acid (50 μmol, Z-A-Y- as listed below) dissolved in 1,2-dichloroethane (3 mL) was added and the array was shaken for 15 min. To each reactor 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine (60 μmol, prepared according to General Procedure B) dissolved in a mixture of acetonitrile (250 μL) and triethylamine (210 μL, 300 μmol) was added and the array was shaken overnight. 1,2-Dichloroethane (1 mL) was added to each reactor and the array was shaken for 2 h. A 0.3 N hydrochloric acid solution (500 μL) was added to each reactor and the array was shaken for 2 h. The lower phase of each reactor was isolated with a pipette and concentrated under reduced pressure. This afforded the following six amides, identified by their MH<sup>+</sup> (LC-MS):

Example	Z-A-Y-	MH <sup>+</sup> (calcd)	MH <sup>+</sup> (found)
14-001		262	262
14-002		248	248
14-003		234	
14-004		302	
14-005		326	
14-006		298	298

**Example 15:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 4-*tert*-butylbenzyl ester oxalate

5



To a solution of 4-*tert*-butylbenzyl alcohol (1.97 g, 12 mmol) in DCM (30 mL) was added pyridine (1.3 mL) and then a solution of 4-nitrophenyl chloroformate (1.57 g, 7.8 mmol) in DCM (20 mL). The mixture was stirred at room temperature for 1 h, concentrated under reduced pressure, and to the residue was added a mixture of 4,5,6,7-tetrahydroimidazo[4,5-c]pyridine (1.5 g, 7 mmol), methanol (3.3 mL), DMF (30 mL), and diisopropylethylamine (4.1 mL). The resulting mixture was stirred at room temperature overnight and concentrated under reduced pressure. The residue was distributed between water and ethyl acetate, phases were separated, and the

15

5390.029-DK,

organic layer was washed with water, dried ( $\text{MgSO}_4$ ), and filtered. A solution of oxalic acid (0.63 g) in ethyl acetate was added to the filtrate and the mixture was allowed to stand for 24 h. Filtration yielded the title compound as colourless solid.

- 5  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  1.28 (s, 9H), 2.65 (m, 2H), 3.69 (t,  $J = 5$  Hz, 2H), 4.46 (s, 2H), 5.08 (s, 2H), 7.31 (d,  $J = 8$  Hz, 2H), 7.39 (d,  $J = 8$  Hz, 2H), 8.13 (m, 1H).

( $\text{C}_{18}\text{H}_{23}\text{N}_3\text{O}_2$ ,  $\text{C}_2\text{H}_2\text{O}_4$ ); calcd. 59.54C 6.25H 10.42N; found 59.66C 6.28H 10.35N.

- 10 The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 4-(*tert*-butyl)benzyl alcohol.

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  1.28 (s, 9H), 2.65 (m, 2H), 3.69 (t,  $J = 5$  Hz, 2H), 4.46 (s, 2H), 5.08 (s, 2H), 7.31 (d,  $J = 8$  Hz, 2H), 7.39 (d,  $J = 8$  Hz, 2H), 8.13 (m, 1H).


15

( $\text{C}_{18}\text{H}_{23}\text{N}_3\text{O}_2$ ,  $\text{C}_2\text{H}_2\text{O}_4$ ); calcd. 59.54C 6.25H 10.42N; found 59.66C 6.28H 10.35N.

Using the same procedure as described for the above example the following compounds were prepared:


20

**Example 16:** 1,4,6,7-Tetrahydro-Imidazo[4,5-*c*]pyridine-5-carboxylic acid benzyl ester



25

**Example 17:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid 2,3-dimethylbenzyl ester oxalate




5

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2,3-dimethylbenzyl alcohol.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  2.19 (s, 3H), 2.27 (s, 3H), 2.63 (m, 2H), 3.70 (t, *J* = 6 Hz, 2H), 4.46 (s, 2H), 5.12 (s, 2H), 7.04–7.18 (m, 3H), 8.15 (s, br, 1H).

10

**Example 18:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid 2-phenylbutyl ester oxalate



15

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2-phenyl-1-butanol oxalate.


M.p. 154–156 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  0.76 (t, *J* = 7 Hz, 3H), 1.53–1.66 (m, 1H), 1.70–1.80 (m, 1H), 2.40–2.63 (m, 2H), 2.82 (m, 1H), 3.48–3.65 (m, 2H), 4.11–4.22 (m, 2H), 4.25–4.40 (m, 2H), 7.18–7.34 (m, 5H), 8.09 (s, 1H).

20

Calcd. 58.60C 5.95H 10.79N; found 58.89C 5.93H 10.63N.

25

**Example 19:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid 3-methoxybenzyl ester oxalate




5

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2-phenyl-1-butanol oxalate.

M.p. 163-165 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.65 (t, *J* = 5 Hz, 2H), 3.69–3.75 (m, 5H), 4.48 (m, 2H), 5.09 (s, 2H), 6.87–6.95 (m, 3H), 7.29 (t, *J* = 8 Hz, 1H), 8.12 (s, 1H).

(C<sub>15</sub>H<sub>17</sub>N<sub>3</sub>O<sub>3</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>); calcd. 54.11C 5.08H 11.14N; found 53.90C 5.04H 11.09N.


**Example 20:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid cyclopentylmethyl ester oxalate



The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and cyclopentylmethanol.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.22 (m, 2H), 1.45–1.61 (m, 4H), 1.62–1.75 (m, 2H), 2.16 (sept, *J* = 7 Hz, 1H), 2.63 (m, 2H), 3.68 (t, *J* = 5 Hz, 2H), 3.92 (d, *J* = 7 Hz, 2H), 4.44 (s, 2H), 8.16 (s, 1H).

**Example 21:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 3-methylbenzyl ester oxalate




5

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 3-methylbenzyl alcohol.

M.p. 169-171 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.31 (s, 3H), 2.65 (m, 2H), 3.70 (m, 2H), 4.47 (s, 2H), 5.08 (s, 2H), 7.11–7.19 (m, 3H), 7.26 (t, *J* = 8 Hz, 1H), 8.15 (s, 1H).

(C<sub>15</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>); calcd. 56.51C 5.30H 11.63N; found 56.74C 5.29H 11.63N.

**Example 22:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-methylbenzyl ester oxalate




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2-methylbenzyl alcohol.

M.p. 158-160 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.31 (s, 3H), 2.64 (s, 2H), 3.69 (t, *J* = 5 Hz, 2H), 4.46 (s, 2H), 5.12 (s, 2H), 7.15–7.26 (m, 3H), 7.31 (d, *J* = 8 Hz, 1H), 8.14 (s, 1H).

Calcd. 56.51C 5.30H 11.63N; found 56.38C 5.27H 11.43N.

**Example 23:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-(2-methoxyphenyl)ethyl ester oxalate



5


The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2-(2-methoxyphenyl)-1-ethanol.

10 M.p. 165-167 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.53–2.63 (m, 2H), 2.88 (t, *J* = 5 Hz, 2H), 3.61 (s, br, 2H), 3.77 (s, 3H), 4.19 (t, *J* = 7 Hz, 2H), 4.39 (s, br, 2H), 6.85 (s, br, 1H), 6.95 (d, *J* = 8 Hz, 1H), 7.10–7.24 (m, 2H), 8.15 (s, 1H).

Calcd. 55.24C 5.41H 10.74N; found 55.39C 5.39H 10.62N.

15

**Example 24:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-(3-methylphenyl)ethyl ester oxalate



20

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2-(3-methylphenyl)-1-ethanol.


25 M.p. 155-157 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.27 (s, 3H), 2.52–2.65 (m, 2H), 2.85 (t, *J* = 5 Hz, 2H), 3.63 (s, br, 2H), 4.21 (t, *J* = 5 Hz, 2H), 4.40 (s, 2H), 7.02 (m, 3H), 7.18 (s, br, 1H), 8.16 (s, 1H).

5390.029-DK,



(C<sub>16</sub>H<sub>19</sub>N<sub>3</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>); calcd. 57.59C 5.64H 11.19N; found 57.33C 5.64H 11.13N.


**Example 25:** 4,5,6,7-Tetrahydro-imidazo[4,5-c]pyridine-5-carboxylic acid adaman-  
tan-1-ylmethyl ester oxalate



The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and  
1-adamantylmethanol.

M.p. 212-214 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.51 (s, 6H), 1.58–1.72 (m, 6H),  
1.94 (s, br, 3H), 2.64 (s, br, 2H), 3.63–3.73 (m, 4H), 4.44 (m, 2H), 8.13 (s, br, 1H).

**Example 26:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid 3,4-  
dimethylbenzyl ester oxalate




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and  
3,4-dimethylbenzyl alcohol.

M.p. 193-195 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.20 (s, 6H), 2.62 (m, 2H), 3.69  
(m, 2H), 4.45 (s, 2H), 5.03 (s, 2H), 7.05–7.14 (m, 3H), 8.12 (s, br, 1H).

Calcd. 57.59C 5.64H 11.19N; found 57.76C 5.60H 10.90N.

**Example 27:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid cyclohex-3-enylmethyl ester oxalate




5

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and cyclohexen-4-ylmethanol.

10  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  1.26 (m, 1H), 1.70–1.80 (m, 2H), 1.82–1.92 (m, 1H), 1.98–2.10 (m, 3H), 2.63 (t,  $J = 5$  Hz, 2H), 3.68 (t,  $J = 5$  Hz, 2H), 3.94 (d,  $J = 7$  Hz, 2H), 4.43 (s, br, 2H), 5.65 (m, 2H), 8.10 (s, br, 1H).

**Example 28:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2,4-dimethylbenzyl ester oxalate

15



The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2,4-dimethylbenzylalcohol.

20

M.p. 176–178 °C.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  2.26 (s, 6H), 2.62 (m, 2H), 3.68 (m, 2H), 4.43 (s, 2H), 5.08 (s, 2H), 6.98 (d,  $J = 8$  Hz, 1H), 7.02 (s, 1H), 7.19 (d,  $J = 8$  Hz, 1H), 8.09 (s, br, 1H).


25

( $\text{C}_{16}\text{H}_{19}\text{N}_3\text{O}_2$ ,  $\text{C}_2\text{H}_2\text{O}_4$ ); calcd. 57.59C 5.64H 11.19N; found 57.51C 5.63H 11.19N.

5390.029-DK,

**Example 29:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-(2-methylphenyl)ethyl ester oxalate

5




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2-(2-methylphenyl)ethanol.

10  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  2.29 (s, 3H), 2.59 (s, br, 2H), 2.91 (m, 2H), 3.63 (s, br, 2H), 4.21 (t,  $J = 5$  Hz, 2H), 4.41 (s, 2H), 7.06–7.20 (m, 4H), 8.15 (s, 1H).

( $\text{C}_{16}\text{H}_{19}\text{N}_3\text{O}_2$ ,  $\text{C}_2\text{H}_2\text{O}_4$ ); calcd. 57.59C 5.64H 11.19N; found 57.44C 5.63H 11.16N.

15 **Example 30:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid naphthalen-1-ylmethyl ester oxalate




20 The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 1-naphthylmethanol.

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  2.52–2.68 (m, 2H), 3.69 (m, 2H), 4.43 (m, 2H), 5.58 (s, 2H), 7.46–7.62 (m, 4H), 7.92–8.00 (m, 2H), 8.05–8.15 (m, 2H).

25

**Example 31:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 4-isopropylbenzyl ester oxalate




5

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 4-isopropylbenzyl alcohol.

M.p. 166-168 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  1.19 (d, *J* = 7 Hz, 6H), 2.64 (m, 2H), 2.89 (sept, *J* = 7 Hz, 1H), 3.70 (t, *J* = 5 Hz, 2H), 4.46 (s, br, 2H), 5.07 (s, 2H), 7.23 (d, *J* = 8 Hz, 2H), 7.29 (d, *J* = 8 Hz, 2H), 8.18 (s, br, 1H).

(C<sub>17</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>); calcd. 58.60C 5.95H 10.79N; found 58.71C 5.97H 10.76N.

**Example 32:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid naphthalen-2-ylmethyl ester oxalate




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2-naphthylmethanol.

M.p. 189-191 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  2.67 (m, 2H), 3.64 (m, 2H), 4.43–4.60 (m, 2H), 5.79 (s, 2H), 7.52–7.56 (m, 3H), 7.89–8.95 (m, 4H), 8.18 (s, br, 1H).

25

Calcd. 60.45C 4.82H 10.57N; found 60.33C 4.82H 10.53N.

**Example 33:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-methylbiphenyl-3-ylmethyl ester oxalate




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2-methylbiphenyl-3-ylmethanol.

10 M.p. 151-153 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.18 (s, 3H), 2.68 (m, 2H), 3.71 (m, 2H), 4.48 (s, br, 2H), 5.19 (s, 2H), 7.18–7.48 (m, 8H), 8.16 (s, br, 1H).

(C<sub>23</sub>H<sub>23</sub>N<sub>3</sub>O<sub>6</sub>, 2/3 H<sub>2</sub>O); calcd. 61.46C 5.46H 9.35N; found 61.62C 5.42H 8.94N.

15 **Example 34:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid 4-ethylbenzyl ester oxalate




20 The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 4-ethylbenzyl alcohol.

M.p. 171-173 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.17 (t, *J* = 7 Hz, 3H), 2.55–2.68 (m, 4H), 3.69 (t, *J* = 5 Hz, 2H), 4.45 (s, br, 2H), 5.08 (s, 2H), 7.20 (d, *J* = 8 Hz, 2H),  
25 7.29 (d, *J* = 8 Hz, 2H), 8.11 (s, br, 1H).

(C<sub>16</sub>H<sub>19</sub>N<sub>3</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>); calcd. 57.59C 5.64H 11.19N; found 57.48C 5.61H 11.20N.

**Example 35:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid thiophen-2-ylmethyl ester oxalate

5




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2-thienylmethanol.

10

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  2.63 (s, br, 2H), 3.68 (s, br, 2H), 4.43 (s, 2H), 5.28 (s, 2H), 7.01 (m, 1H), 7.17 (d, *J* = 3 Hz, 1H), 7.54 (d, *J* = 5 Hz, 1H), 8.10 (m, 1H).

**Example 36:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-thiophen-2-ylethyl ester oxalate

15



The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2-(2-thienyl)-1-ethanol.

20


M.p. 173-175 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  2.62 (s, br, 2H), 3.13 (t, *J* = 7 Hz, 2H), 3.68 (t, *J* = 5 Hz, 2H), 4.21 (t, *J* = 7 Hz, 2H), 4.43 (s, br, 2H), 6.89–6.99 (m, 2H), 7.34 (s, br, 1H), 8.11 (s, br, 1H).

25

(C<sub>13</sub>H<sub>15</sub>N<sub>3</sub>O<sub>2</sub>S); calcd. 49.04C 4.66H 11.44N; found 49.12C 4.62H 11.43N.

5390.029-DK,

**Example 37:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-thiophen-3-ylethyl ester oxalate



5


The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2-(3-thienyl)-1-ethanol.

- 10 M.p. 193-194 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.59 (s, br, 2H), 2.93 (t, *J* = 7 Hz, 2H), 3.63 (s, br, 2H), 4.21 (t, *J* = 7 Hz, 2H), 4.40 (s, 2H), 7.03 (s, br, 1H), 7.24 (s, br, 1H), 7.46 (s, br, 1H), 8.08 (s, br, 1H).

(C<sub>13</sub>H<sub>15</sub>N<sub>3</sub>O<sub>2</sub>S); calcd. 49.04C 4.66H 11.44N; found 49.12C 4.61H 11.40N.

15

**Example 38:** 4,5,6,7-Tetrahydro-imidazo[4,5-*c*]pyridine-5-carboxylic acid 4-methoxy-2,3-dimethylbenzyl ester oxalate




20

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2,3-dimethyl-4-methoxybenzyl alcohol.

- 25 M.p. 161-162 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.09 (s, 3H), 2.20 (s, 3H), 2.62 (s, br, 2H), 3.65 (s, br, 2H), 3.77 (s, 3H), 4.42 (s, br, 2H), 5.08 (s, 2H), 6.78 (d, *J* = 8 Hz, 1H), 7.15 (d, *J* = 8 Hz, 1H), 8.18 (s, br, 1H).

5390.029-DK,


**Example 39:** 4-Trifluoromethyl-4,5,6,7-tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid benzyl ester oxalate



The compound was prepared from 4-trifluoromethyl-4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and benzyl alcohol.

- 10 M.p. 169-171 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  2.65–2.81 (m, 2H), 2.99 (s, br, 2H), 4.42 (q, *J* = 8 Hz, 1H), 5.42 (s, 2H), 7.36–7.45 (m, 3H), 7.51 (d, *J* = 8 Hz, 2H), 8.21 (s, 1H). 1H).

**Example 40:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid 2-cyclohexylethyl ester oxalate



20 The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2-cyclohexyl-1-ethanol.


M.p. 174-176 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  0.83–0.95 (m, 2H), 1.05–1.26 (m, 3H), 1.32 (m, 1H), 1.48 (q, *J* = 7 Hz, 2H), 1.55–1.71 (m, 5H), 2.64 (m, 2H), 3.67 (t, *J* = 5 Hz, 2H), 4.06 (t, *J* = 7 Hz, 2H), 4.44 (s, 2H), 8.22 (s, br, 1H).

(C<sub>15</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub>); calcd. 55.58C 6.86H 11.40N; found 55.70C 6.90H 11.32N.

5390.029-DK,



**Example 41:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid cyclohexylmethyl ester oxalate




5

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and cyclohexylmethanol.

M.p. 176-178 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 0.89–1.03 (m, 2H), 1.06–1.29 (m, 3H), 1.55–1.72 (m, 6H), 2.65 (m, 2H), 3.68 (m, 2H), 3.86 (d, *J* = 7 Hz, 2H), 4.44 (s, br, 2H), 8.20 (s, br, 1H).

(C<sub>14</sub>H<sub>21</sub>N<sub>3</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>) calcd. 54.38C 6.56H 11.89N; found 54.56C 6.64H 11.77N.

**Example 42:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-chlorobenzyl ester oxalate




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and 2-chlorobenzyl alcohol.

M.p. 173-175 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.65 (m, 2H), 3.71 (s, br, 2H), 4.48 (m, 2H), 5.19 (s, 2H), 7.38 (m, 2H), 7.50 (m, 2H), 8.09 (s, br, 1H).

25

(C<sub>14</sub>H<sub>14</sub>ClN<sub>3</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>); calcd. 50.34C 4.22H 11.01N; found 50.21C 4.19H 10.81N.

**Example 43:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid 2-trifluoromethylbenzyl ester oxalate




5

The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 2-(trifluoromethyl)benzyl alcohol.

M.p. 152-156 °C (EtOAc). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.67 (m, 2H), 3.71 (m, 2H), 4.48 (s, br, 2H), 5.28 (s, 2H), 7.58 (t, *J* = 8 Hz, 1H), 7.65–7.81 (m, 3H), 8.14 (m, 1H).

(C<sub>15</sub>H<sub>14</sub>F<sub>3</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>); calcd. 49.16C 3.88H 10.12N; found 49.21C 3.86H 10.11N.

**Example 44:** 4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine-5-carboxylic acid 1-phenylethyl ester oxalate




The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 1-phenylethanol.

M.p. 162 °C (decomposes; EtOAc/MeOH). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 1.49 (d, *J* = 7 Hz, 3H), 2.62 (s, br, 2H), 3.62–3.80 (m, 2H), 4.38–4.60 (m, 2H), 5.74 (q, *J* = 7 Hz, 1H), 7.25–7.40 (m, 5H), 8.06 (s, br, 1H).

Calcd. 56.51C 5.30H 11.63N; found 56.46C 5.28H 11.48N.

**Example 45:** (4-Benzylpiperidin-1-yl)-(4,5,6,7-tetrahydroimidazo[4,5-c]pyridin-5-yl)-methanone oxalate



The compound was prepared from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine and 4-benzylpiperidine according to the following, general procedure for the preparation of ureas:

10 A solution of 4-nitrophenyl chloroformate (1.04 g, 5.2 mmol) in DCM (10 mL) was dropwise added to a solution of 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine (1.0 g, 4.7 mmol) in methanol (14 mL) and triethylamine (1.9 mL). The resulting mixture was stirred at 20 °C for 1 h, concentrated under reduced pressure, and the residue was

15 distributed between DCM (40 mL) and water (40 mL). After separation of phases the aqueous phase was extracted with DCM (1 x 40 mL), and the combined organic phases were dried (MgSO<sub>4</sub>) and concentrated. The residue was dissolved in DMF (20 mL), 4-benzylpiperidine (1.65 mL, 9.39 mmol) was added, and the resulting mixture was stirred at 80 °C for 17 h. The mixture was concentrated under reduced

20 pressure, and the residue was redissolved in ethyl acetate. After washing with water the organic phase was dried (MgSO<sub>4</sub>) and filtered. To the filtrate was added a solution of oxalic acid (0.44 g, 4.9 mmol) in ethyl acetate, and the resulting precipitate is filtered off. The resulting solid was distributed between an aqueous solution of NaHCO<sub>3</sub> and ethyl acetate, and the organic layer was purified by column chromatogra-


25 phy. The purified compound was again precipitated as oxalate from ethyl acetate to yield 0.58 g (30%) of the title oxalate as colorless solid, which does not melt but decomposes upon heating.

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  1.06–1.20 (m, 2H), 1.52–1.60 (m, 2H), 1.61–1.75 (m, 1H), 2.52 (m, 2H), 2.65–2.78 (m, 4H), 3.42 (t,  $J = 5$  Hz, 2H), 3.55–3.63 (m, 2H), 4.23 (s, 2H), 7.16–7.31 (m, 5H), 8.40 (s, 1H).

- 5 Anal. calcd. for  $\text{C}_{19}\text{H}_{24}\text{N}_4\text{O}$ , 1.5  $\text{C}_2\text{H}_2\text{O}_4$  (459.49): 57.51C 5.92H 12.19N; found: 57.67C 6.20H 12.25N.

**Example 46:** 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine-5-carboxylic acid N-methyl-N-benzylamide oxalate

10



The compound was prepared in the same way as disclosed for example 45 from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and benzylmethylamine.


15

M.p. 164–166 °C (EtOAc/MeOH).  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  2.71 (m, 5H), 3.49 (t,  $J = 5$  Hz, 2H), 4.23 (s, 2H), 4.36 (s, 2H), 7.23–7.39 (m, 5H), 8.19 (s, 1H).

Calcd. 56.66C 5.59H 15.55N; found 56.75C 5.61H 15.35N.

20

**Example 47:** (3,4-Dihydro-1*H*-isoquinolin-2-yl)(4,5,6,7-tetrahydroimidazo[4,5-c]pyridin-5-yl)methanone




25

The compound was prepared in the same way as disclosed for example 45 from 4,5,6,7-tetrahydro-1*H*-imidazo[4,5-c]pyridine and benzylmethylamine.

M.p. 102-104 °C (EtOAc). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 2.66 (s, br, 2H), 2.84 (t, *J* = 5 Hz, 2H), 3.46 (m, 4H), 4.21 (s, br, 2H), 4.39 (s, 2H), 7.16 (s, 4H), 7.47 (s, 1H), 11.80 (s, 1H).

5

**Example 48:** 3,3-Dimethyl-1-(1,4,6,7-tetrahydroimidazo[4,5-*c*]pyridin-5-yl)-butan-1-one



10

At 0 °C, EDC (0.45 g, 2.4 mmol) was added to a solution of tert-butylacetic acid (0.30 mL, 2.3 mmol), and 1-hydroxy-7-azabenzotriazole (0.32 g, 2.4 mmol) in dichloromethane (30 mL). The reaction mixture was stirred for 20 min at 0 °C. 4,5,6,7-tetrahydroimidazo[4,5-*c*]pyridine dihydrochloride (0.50 g, 2.4 mmol) was added. Ethyldiisopropylamine (0.40 mL, 2.4 mmol) was added. The reaction mixture was stirred for 16 h at room temperature. The reaction mixture was diluted with ethyl acetate (100 mL) and washed with 10% aqueous sodium hydrogen sulfate solution (100 mL). A 1 N solution of sodium hydroxide was added to the aqueous solution until pH 12 was obtained. It was extracted with ethyl acetate (2 x 100 mL). These organic extracts were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent to give 125 mg of the title compound.

15


20

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 2 rotamers): δ 1.03 and 1.10 (both s, together 9 H); 2.35 and 2.40 (both s, together 2 H); 2.68 and 2.78 (both t, together 2 H); 3.80 and 3.95 (both t, together 2 H); 4.55 and 4.70 (both s, together 2 H); 7.53 (s, 1 H).

For biological testing it was transferred into its acetate salt by lyophilization with 0.5 M acetic acid (40 mL).

**Example 49:** 3-Phenyl-1-(1,4,6,7-tetrahydroimidazo[4,5-c]pyridin-5-yl)propynone

5



1-Hydroxy-7-azabenzotriazole (0.32 g, 2.4 mmol) was added to a solution of phenylpropionic acid (0.34 g, 2.4 mmol) in dichloromethane (30 mL). The solution was cooled to 0 °C. EDC (0.45 g, 2.4 mmol) was added. The reaction mixture was stirred at 0 °C for 20 min. 4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine dihydrochloride (0.50 g, 2.3 mmol) was added. Ethyldiisopropylamine (0.40 mL, 2.3 mmol) was added. The reaction mixture was stirred for 16 h at room temperature. It was diluted with ethyl acetate (100 mL) and washed with 10% aqueous sodium hydrogensulfate solution (100 mL). The aqueous phase was extracted with ethyl acetate (3 x 60 mL): It was added a 1 N sodium hydroxide solution until pH 12 was obtained. It was extracted with ethyl acetate (3 x 90 mL). These extracts were combined and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 50 mg of the title compound.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 2 rotamers): δ 2.75 and 2.85 (both t, together 2 H); 4.03 and 4.15 (both t, together 2 H); 4.75 and 4.92 (both s, together 1 H); 7.10 - 7.70 (m, 6 H).


The title compound was transferred into its oxalate salt by dissolving in ethyl acetate (20 mL) and addition of a solution of oxalic acid in ethyl acetate (20 mL). The precipitation was collected and washed with ethyl acetate (10 mL). It was dried in vacuo.

HPLC :  $R_t$  = 6.922 min (Method h8).

LC-MS calcd. for  $MH^+$ : 252 found: 252.

5

**Example 50:** 2E-3-(4-*tert*-Butylphenyl)-1-(1,4,6,7-tetrahydroimidazo[4,5-*c*]pyridin-5-yl)propenone



10

At 0 °C, EDC (0.45 g, 2.4 mmol) was added to a solution of 3-(4-*tert*-butylphenyl)-acrylic acid (0.48 g, 2.4 mmol) and 1-hydroxy-7-azabenzotriazole (0.32 g, 2.4 mmol) in dichloromethane (30 mL). The reaction mixture was stirred for 20 min at 0 °C.

4,5,6,7-Tetrahydroimidazo[4,5-*c*]pyridine dihydrochloride (0.50 g, 2.3 mmol) was added. Ethyldiisopropylamine (0.40 mL, 2.3 mmol) was added. The reaction mixture was stirred for 16 h at room temperature. It was diluted with ethyl acetate (100 mL) and washed with a saturated aqueous solution of sodium hydrogencarbonate (100 mL). The aqueous phase was extracted with ethyl acetate (2 x 60 mL). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 119 mg of the title compound.

15

20

$^1H$  NMR ( $CDCl_3$ ):  $\delta$  1.32 (s, 9 H); 2.75 (br, 2 H); 3.95 (br, 2 H); 4.75 (br, 2 H); 6.95 (d, 1 H); 7.35 - 7.55 (m, 5 H); 7.65 (d, 1 H).

25

For biological testing, the title compound was transferred into its oxalate salt: The title compound was dissolved in ethyl acetate (15 mL). A solution of oxalic acid 5390.029-DK,

(58 mg, 0.64 mmol) in ethyl acetate (15 mL) was added. The precipitation was collected, washed with ethyl acetate (10 mL) and dried in vacuo.


HPLC:  $R_t$  9.55 min (Method h8).

5

LC-MS: calcd. for  $MH^+$ : 310; found 310.

**Example 51:** 2-Bicyclo[2.2.1]hept-2-yl-1-(1,4,6,7-tetrahydroimidazo[4,5-c]pyridin-5-yl)ethanone

10



At 0 °C, EDC (0.45 g, 2.4 mmol) was added to a solution of 1-hydroxy-7-azabenzotriazole (0.32 g, 2.4 mmol) and 2-norbornaneacetic acid (0.338 mL, 2.4 mmol) in dichloromethane (30 mL). The reaction mixture was stirred for 20 min at 0 °C.

15

4,5,6,7-Tetrahydroimidazo[4,5-c]pyridine dihydrochloride (0.50 g, 2.3 mmol) was added. Ethyldiisopropylamine (0.40 mL, 2.3 mmol) was added. The reaction mixture was stirred for 16 h at room temperature. It was diluted with ethyl acetate (100 mL) and washed with a saturated aqueous solution of sodium hydrogencarbonate (100 mL). The aqueous solution was extracted with ethyl acetate (2 x 60 mL). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 120 mg of the title compound.

20

25

$^1H$  NMR ( $CDCl_3$ , 2 rotamers):  $\delta$  1.00 - 2.50 (m, 13 H); 2.65 and 2.75 (both t, together 2 H); 3.75 and 3.90 (both t, together 2 H); 4.55 and 4.65 (both s, together 2 H); 7.53 (s, 1 H).

5390.029-DK,



For biological testing, the title compound was transferred into its oxalate salt: The title compound was dissolved in ethyl acetate (15 mL). A solution of oxalic acid (42 mg, 0.47 mmol) in ethyl acetate (15 mL) was added. The precipitation was collected, washed with ethyl acetate (10 mL) and dried in vacuo.

HPLC:  $R_t = 7.59$  min (Method h8).


LC-MS: calcd. for  $MH^+$ : 260; found 260.

$C_{15}H_{21}N_3O \cdot C_2H_2O_4$  (349.39)










calcd: 58.44C 6.64H 12.03N;










found: 58.72C 6.70H 11.99N.









- Furthermore, the following preferred compounds according to the invention may be prepared according to the general procedures set forth in the foregoing description:


















Example No	-X-Y-A-Z
52	
53	










54	 <chem>C=C(C)N(Cc1ccccc1)N#C</chem>
55	 <chem>C=C(C)N(Cc1ccc(Cl)cc1)N#C</chem>
56	 <chem>C=C(C)N(Cc1ccc(cc1)C(=O)C2CC2)N#C</chem>
57	 <chem>C=C(C)N(CCCc1ccccc1)N#C</chem>
58	 <chem>C=C(C)N(CCCc1ccc(Cl)cc1)N#C</chem>
59	 <chem>C=C(C)N(CCCc1ccc(cc1)C(=O)C2CC2)N#C</chem>
60	 <chem>C=C(C)N(CCCc1ccccc1)N#C</chem>
61	 <chem>C=C(C)N(CCCc1ccc(Cl)cc1)N#C</chem>
62	 <chem>C=C(C)N(CCCc1ccc(cc1)C(C)(C)C)N#C</chem>












63	
64	
65	
66	
67	
68	
69	
70	
71	

72	
73	
74	
75	
76	
77	
78	
79	






80	
81	
82	
83	
84	
85	
86	
87	

88	
89	
90	
91	
92	
93	
94	

95	
96	
97	
98	
99	
100	
101	
102	
103	

104	
105	
106	
107	
108	
109	
110	
111	
112	
113	
114	



115	
116	
117	
118	
119	

## PHARMACOLOGICAL METHODS

### In vitro binding assay

- 5 The ability of the compounds to interact with the histamine H3 receptor was determined by an *in vitro* binding assay. Rat cerebral cortex was homogenized in ice cold K-Hepes, 5 mM MgCl<sub>2</sub> pH 7.1 buffer. After two differential centrifugations the last pellet was resuspended in fresh Hepes buffer containing 1 mg/mL Bacitracin. Aliquots of the membrane suspension (400 mg/mL) were incubated for 60 min at
  - 10 25 °C with 30 pM [<sup>125</sup>I]-iodoproxifan, a known histamine H3 receptor antagonist, and the test compound at various concentrations. The incubation was stopped by dilution with ice-cold medium, followed by rapid filtration through Whatman GF/B filters pretreated for 1 h with 0.5% polyethyleneimine. The radioactivity retained on the filters was counted using a Cobra II auto gamma counter. The radioactivity of the
- 5390.029-DK,

filters was indirectly proportional to the binding affinity of the tested compound. The results were analyzed by nonlinear regression analysis.

When tested the present compounds showed a high binding affinity to the histamine  
5 H3 receptor.

Preferably, the compounds according to the invention have an  $IC_{50}$  value as determined by the assay of less than 1  $\mu$ M, more preferred of less than 500 nM and even more preferred of less than 100 nM.

10

Furthermore, in a similar way binding assays were carried out in order to determine the ability of the present compounds to interact with the histamine H1 receptor (reference compound [ $^{125}$ I]-pyrilamine) and the histamine H2 receptor (reference compound [ $^{125}$ I]-aminopotentidine), respectively. These assays showed that the present compounds do not show a high affinity for these receptors and hence are very  
15 specific for the histamine H3 receptor.

#### The open cage Schedule-fed rat model

The ability of the present compounds to reduce weight was determined using the *in*  
20 *vivo* open cage Schedule-fed rat model.

Sprague-Dawley (SD) male rats of an age of about 1½ to 2 months and a weight of about 250 g were habituated to the presence of food (Altromin pelleted rat chow) in their home cage only during three hours in the morning from 9 to 12 a.m. all days a  
25 week. Water was present ad libitum. As the consumption of food stabilised after 7 to 9 days, the animals were ready for use.

The animals were tested twice a week. During the test sessions, the test compound was administered intraperitoneally 30 minutes before the start of the sessions. One  
30 group of 9 animals was administered the test compound at a dose of 15 mg/kg and another group of 11 animals was administered the test compound at a dose of 30

mg/kg. A control group of 11 animals was administered the vehicle consisting of NaCl 0.9% and Chremophor 5%. Food and water intake were monitored at 1, 2 and 3 h post administration.


- 5 During the test period the animals were weighed weekly and if necessary extra food was given in order to ensure that the weight gain was 3 to 5 g per week corresponding to the normal weight gain for SD male rats at this age.

- 10 Any side effects could rapidly be discovered (barrel-rolling, bushy fur etc.) since the animals were kept in transparent plastic cages to enable continuous monitoring.

When tested the compounds significantly and dose-dependently inhibits food intake.

# CLAIMS

1. A compound of the general formula I



wherein

R<sup>1</sup> is hydrogen or a functional group which can be converted to hydrogen *in vivo*;

R<sup>2</sup> is hydrogen, C<sub>1-6</sub>-alkyl, halogen, cyano, trifluoromethyl, hydroxy or -NR<sup>7</sup>R<sup>8</sup>,

wherein R<sup>7</sup> and R<sup>8</sup> independently are

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

5 C<sub>1-6</sub>-alkylsulfonyl optionally substituted with  
C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

10 R<sup>7</sup> and R<sup>8</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;  
15

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> independently are

hydrogen, carboxy, C<sub>1-6</sub>-alkoxycarbonyl, cyano, trifluoromethyl, halogen,

20

C<sub>3-8</sub>-cycloalkyl optionally substituted with  
C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

25

C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with  
C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, cyano, halogen, trifluoromethyl, carboxy, C<sub>1-6</sub>-alkoxycarbonyl,

30 C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

5 aryl optionally substituted with

halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, heteroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

10 -CO-NR<sup>9</sup>R<sup>10</sup>,

wherein R<sup>9</sup> and R<sup>10</sup> independently are

hydrogen,

15

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

20 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

25 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

5 R<sup>9</sup> and R<sup>10</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl,  
10 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, and/or R<sup>5</sup> and R<sup>6</sup> together with the carbon atom to which they are connected, form a 3 to 8 mem-  
15 bered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or hetero-aryl-amino;

20 m, n, p and q independently are 0, 1 or 2;

X is a valence bond, -CH<sub>2</sub>-, -C(=O)-, -C(=S)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)-, -C[=C(CN)<sub>2</sub>]-, -C(=CH-CN)-, -C(=NR<sup>11</sup>)- or -C(=N-S(=O)<sub>2</sub>R<sup>11a</sup>)-,

25 wherein R<sup>11</sup> is

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with  
30 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

5 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

10

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

15

R<sup>11a</sup> is

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

20

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

25

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

30

Y is a valence bond, -O- or -N(R<sup>12</sup>)-,



wherein R<sup>12</sup> is

hydrogen,

5

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

10

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroaryl-amino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

15

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

20

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

A is a valence bond, C<sub>1-8</sub>-alkylene, C<sub>2-8</sub>-alkenylene, C<sub>2-8</sub>-alkynylene, C<sub>3-8</sub>-cyclo-alkylene or phenylene, or

25

when Y is -N(R<sup>12</sup>)-, A, together with R<sup>12</sup> and the nitrogen atom to which they are connected, may form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring system optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl,

30

heteroaryl, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and

Z is -R<sup>13</sup>, -OR<sup>13</sup>, -SR<sup>13</sup>, -NR<sup>13</sup>R<sup>14</sup>, -CHR<sup>13</sup>R<sup>14</sup>, -CR<sup>13</sup>R<sup>14</sup>R<sup>15</sup> or =CR<sup>13</sup>R<sup>14</sup>,

5

wherein R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> independently are

hydrogen,

10

C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with aryl, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, heteroaryl or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

15

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino, halogen, cyano or trifluoromethyl,

20

aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, C<sub>3-8</sub>-cycloalkanecarbonyl, hydroxy, amino, halogen, cyano or trifluoromethyl, where

25

R<sup>13</sup> and R<sup>14</sup> or R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup>, when they do not represent hydrogen, may be joined by one or more bridging linkers such as a valence bond, C<sub>1-4</sub>-alkylene, C<sub>2-4</sub>-alkenylene, -O-, -S-, -N(R<sup>16</sup>)-, -C(=O)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(R<sup>16</sup>R<sup>17</sup>)-, phenylene, biphenylene, -O-C<sub>1-4</sub>-alkylene, -S-C<sub>1-4</sub>-alkylene, -N(R<sup>16</sup>)-C<sub>1-4</sub>-alkylene,

30

-N=C<sub>1-4</sub>-alkylene, -O-C<sub>2-4</sub>-alkenylene, -S-C<sub>2-4</sub>-alkenylene or -N(R<sup>16</sup>)-C<sub>2-4</sub>-alkenylene, to form a mono-, bi- or polycyclic ring system,

wherein R<sup>16</sup> and R<sup>17</sup> independently are

5

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

10

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino,

15

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroaryl-amino,

20

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

25

R<sup>16</sup> and R<sup>17</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl,

30

C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

with the provisos that

5

when X is -CS-,  $R^1 = R^2 = R^5 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ , the group -Y-A-Z must not start with the radical -NH-;

when the group -X-Y-A-Z starts with the radical -CH<sub>2</sub>-,  $R^1 = R^2 = R^6 = \text{hydrogen}$ ,

10

$m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy or aminocarbonyl;

when X is -CO-, the group -Y-A-Z starts with the radical -NH-,  $R^1 = R^2 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ , the remainder of the group -Y-A-Z must not be hydrogen, unsubstituted or C<sub>1-6</sub>-alkoxy substituted phenyl, unsubstituted C<sub>3-8</sub>-cycloalkyl or

15

unsubstituted C<sub>1-6</sub>-alkyl;

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl,  $R^1 = R^2 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy, aminocarbonyl or 4-phenyl-piperazin-1-ylcarbonyl;

20

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl,  $R^1 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $R^2 = \text{butyl}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be methoxycarbonyl;

when X is -CO-, Y is -O-, A is -CH<sub>2</sub>-, Z is phenyl,  $R^1 = R^2 = R^4 = R^5 = R^6 = \text{hydrogen}$ ,

25

$m = n = p = 0$  and  $q = 1$ ,  $R^3$  must not be hydrogen, ethyl, isopropyl or phenyl;

when X is -CO-, Y is -O-, A is a valence bond, Z is *tert*-butyl,  $R^1 = R^2 = R^3 = R^4 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy;

30

when X is -CO-, Y is -O-, A is a valence bond, Z is *tert*-butyl,  $R^1 = R^2 = R^4 = R^5 = R^6 = \text{hydrogen}$ ,  $m = n = p = 0$  and  $q = 1$ ,  $R^3$  must not be 4-cyanophenyl;

when X is -CO-, the group -Y-A-Z starts with the radical -O-,  $R^1 = R^2 = R^6 =$  hydrogen,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy, aminocarbonyl or hydrogen;

- 5 when -X is -CO-, the group -Y-A-Z starts with the radical -CH<,  $R^1 = R^2 = R^3 = R^4 = R^6 =$  hydrogen,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be hydroxymethyl,  $C_{1-6}$ -alkoxy-carbonyl or carboxy; and

when X is -CO-, the group -Y-A-Z is 4-methoxyphenyl,  $R^1 = R^2 = R^3 = R^4 = R^6 =$  hydrogen,  $m = n = p = 0$  and  $q = 1$ ,  $R^5$  must not be carboxy;

as well as any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof.

- 15 2. A compound according to claim 1 wherein  $R^1 = R^2 =$  hydrogen;  $m = n = p = 0$ ; and  $q = 1$ .

3. A compound according to claim 1 or 2 wherein X is -C(=O)-; Y is a valence bond; A is a valence bond or  $C_{1-8}$ -alkylene; and Z is  $-R^{13}$ ,  $-CHR^{13}R^{14}$  or  $-NHR^{13}R^{14}$ ; wherein  $R^{13}$  and  $R^{14}$  are as defined in claim 1.

4. A compound according to claim 3 wherein Z is  $-R^{13}$  in which  $R^{13}$  is

aryl,  $C_{3-15}$ -cycloalkyl, aroyl or heteroaryl, which are optionally substituted with  
 25 aryl- $C_{1-6}$ -alkyl, heteroaryl- $C_{1-6}$ -alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl,  $C_{1-6}$ -alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino,  $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxy,  $C_{1-6}$ -alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl.

- 30 5. A compound according to claim 3 wherein Z is  $-CHR^{13}R^{14}$  in which  $R^{13}$  and  $R^{14}$  independently are

hydrogen, or

aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with

- 5        aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl.

- 10    6. A compound according to claim 3 wherein Z is -NR<sup>13</sup>R<sup>14</sup> in which R<sup>13</sup> and R<sup>14</sup> independently represent

hydrogen,

- 15    aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl, or

20

R<sup>13</sup> and R<sup>14</sup> each independently represent aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are joined with a C<sub>1-4</sub>-alkylene group to form a polycyclic ring system.

- 25    7. A compound according to claim 1 or 2 wherein X is -C(=O)-; Y is -NH-; A is a valence bond, C<sub>1-8</sub>-alkylene or C<sub>3-8</sub>-cycloalkylene; and

Z is -R<sup>13</sup> in which R<sup>13</sup> is

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, heteroaryl or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

5 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroaryl-sulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryl-oxy, acylamino, hydroxy, amino, halogen, cyano or trifluoromethyl,

aryl, C<sub>3-15</sub>-cycloalkyl or heteroaryl, which are optionally substituted with

10 aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroaryl-sulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl, or

15 Z is -CHR<sup>13</sup>R<sup>14</sup> in which R<sup>13</sup> and R<sup>14</sup> independently are

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

20 C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroaryl-sulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino, halogen, cyano or trifluoromethyl,

25 aryl, C<sub>3-15</sub>-cycloalkyl or heteroaryl, which are optionally substituted with

aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroaryl-amino, aroyl, heteroaroyl, arylsulfonyl, heteroaryl-sulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano or trifluoromethyl.

30

8. A compound according to any one of the preceding claims wherein R<sup>3</sup> and R<sup>4</sup> independently are

hydrogen;

5

C<sub>3-8</sub>-cycloalkyl optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

10

C<sub>1-6</sub>-alkyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

15

aryl optionally substituted with

halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, heteroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup> wherein R<sup>9</sup> and R<sup>10</sup> are as defined in claim 1, or

20

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, form a C<sub>3-8</sub>-cycloalkyl ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino.

25

9. A compound according to any one of the claims 1 to 8 wherein R<sup>5</sup> and R<sup>6</sup> are both hydrogen.

30

10. A compound according to any one of the claims 1 to 9 wherein R<sup>3</sup> and R<sup>4</sup> are both hydrogen or are both C<sub>1-6</sub>-alkyl, or R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to



which they are connected, form a C<sub>3-8</sub>-cycloalkyl ring, or one of R<sup>3</sup> and R<sup>4</sup> is hydrogen while the other is C<sub>3-8</sub>-cycloalkyl substituted C<sub>1-6</sub>-alkyl.

11. A compound according to any one of the claims 1 to 10 wherein R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are hydrogen.

12. A compound according to claim 1 or 2 wherein X is -C(O)- and Y is -O-.

13. A compound according to claim 12 wherein A is C<sub>1-6</sub>-alkylene or a valence bond.

14. A compound according to claim 13 wherein A is methylene or ethylene.

15. A compound according to claim 13 or 14 wherein Z is -R<sup>13</sup>, -CHR<sup>13</sup>R<sup>14</sup> or -CR<sup>13</sup>R<sup>14</sup>R<sup>15</sup> wherein R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> are as defined in claim 1.

16. A compound according to claim 15 wherein Z is aryl, heteroaryl or C<sub>3-15</sub>-cycloalkyl which are optionally substituted as defined in claim 1.

17. A compound according to claim 16 wherein Z is phenyl, naphthyl, thienyl, cyclopentyl, cyclohexyl or cyclohexenyl which are optionally substituted as defined in claim 1.

18. A compound according to claim 17 wherein Z is phenyl, naphthyl, thienyl, cyclopentyl, cyclohexyl or cyclohexenyl.

19. A compound according to claim 17 wherein Z is phenyl which is substituted with one to three substituents selected from C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, halogen, phenyl and trifluoromethyl.

20. A compound according to claim 15 wherein Z is -CHR<sup>13</sup>R<sup>14</sup> in which R<sup>13</sup> is C<sub>1-6</sub>-alkyl and R<sup>14</sup> is phenyl.

21. A compound according to claim 15 wherein Z is a polycarbocyclic ring system.

22. A compound according to claim 20 wherein Z is adamantyl.

5

23. A compound according to any one of the claims 12 to 22 wherein R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are hydrogen.

24. A compound according to claim 1 wherein m = n = p = 0 and q = 1; R<sup>1</sup> = R<sup>2</sup> = hydrogen; R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are hydrogen; X is -C(O)-; Y is -N(R<sup>12</sup>)- wherein R<sup>12</sup> and A, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring system optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is hydrogen.

10

15

25. A compound according to claim 1 wherein m = n = p = 0 and q = 1; R<sup>1</sup> = R<sup>2</sup> = hydrogen; R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are hydrogen; X is -C(O)-; Y is -N(R<sup>12</sup>)- wherein R<sup>12</sup> is C<sub>1-6</sub>-alkyl; A is C<sub>1-8</sub>-alkylene; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is aryl optionally substituted as defined in claim 1.

20

26. A compound according to claim 1 wherein m = n = p = 0 and q = 1; R<sup>1</sup> = R<sup>2</sup> = hydrogen; R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are hydrogen; X is -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)- or -C(=N-S(=O)<sub>2</sub>R<sup>11a</sup>)- wherein R<sup>11a</sup> is C<sub>1-6</sub>-alkyl or phenyl substituted with C<sub>1-6</sub>-alkyl; Y is -NH-; A is C<sub>1-8</sub>-alkylene; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is aryl or C<sub>3-15</sub>-cycloalkyl optionally substituted as defined in claim 1.

25

27. A compound according to claim 1 wherein m = n = p = 0 and q = 1; R<sup>1</sup> = R<sup>2</sup> = hydrogen; R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are hydrogen; X is -S(=O)<sub>2</sub>-; Y is a valence bond; A is a

30

valence bond or C<sub>1-8</sub>-alkylene; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is aryl or C<sub>3-15</sub>-cycloalkyl optionally substituted as defined in claim 1.

5 28. A compound according to claim 1 wherein m = n = p = 0 and q = 1; R<sup>1</sup> = R<sup>2</sup> = hydrogen; R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are hydrogen; X is -C(=O)-; Y is -NH-; A is a valence bond or C<sub>1-8</sub>-alkylene; and Z is -R<sup>13</sup> wherein R<sup>13</sup> is aryl or C<sub>3-15</sub>-cycloalkyl optionally substituted as defined in claim 1.

10 29. A compound according to any one of the claims 26 to 28 wherein R<sup>13</sup> is phenyl or cyclohexyl which are optionally substituted with halogen, C<sub>1-6</sub>-alkyl or C<sub>3-8</sub>-cycloalkanecarbonyl.


15 30. A compound according to any one of the claims 26 to 29 wherein A is methylene, ethylene or propylene.

31. A compound according to any one of the claims 1 to 30 for use as a medicament.

20 32. A pharmaceutical composition comprising, as an active ingredient, at least one compound according to any one of the claims 1 to 30 together with one or more pharmaceutically acceptable carriers or diluents.

25 33. A pharmaceutical composition according to claim 32 in unit dosage form, comprising from about 0.05 mg to about 1000 mg, preferably from about 0.1 mg to about 500 mg and especially preferred from about 0.5 mg to about 200 mg of the compound according to any one of the claims 1 to 30.

34. Use of a compound of the general formula I'



5 wherein

R<sup>1</sup> is hydrogen or a functional group which can be converted to hydrogen *in vivo*;

R<sup>2</sup> is hydrogen, C<sub>1-6</sub>-alkyl, halogen, cyano, trifluoromethyl, hydroxy or -NR<sup>7</sup>R<sup>8</sup>,

10

wherein R<sup>7</sup> and R<sup>8</sup> independently are

hydrogen,

15

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

20

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

5 C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, or

10 R<sup>7</sup> and R<sup>8</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> independently are

hydrogen, carboxy, C<sub>1-6</sub>-alkoxycarbonyl, cyano, trifluoromethyl, halogen,

20 C<sub>3-8</sub>-cycloalkyl optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

25 C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with

C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, cyano, halogen, trifluoromethyl, carboxy, C<sub>1-6</sub>-alkoxycarbonyl,

30 C<sub>3-8</sub>-cycloalkyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

5 aryl optionally substituted with

halogen, cyano, nitro, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, hydroxy, trifluoromethyl, aryl, heteroaryl, aryloxy, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroarylamino or -CO-NR<sup>9</sup>R<sup>10</sup>,

10 -CO-NR<sup>9</sup>R<sup>10</sup>,

wherein R<sup>9</sup> and R<sup>10</sup> independently are

hydrogen,

15

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-  
20 lamino or heteroarylamino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

25

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

5 R<sup>9</sup> and R<sup>10</sup>, together with the nitrogen atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl,  
10 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino, or

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are connected, and/or R<sup>5</sup> and R<sup>6</sup> together with the carbon atom to which they are connected, form a 3 to 8 membered, saturated or unsaturated, carbocyclic or heterocyclic ring optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;  
15

20 m, n, p and q independently are 0, 1 or 2;

X is a valence bond, -CH<sub>2</sub>-, -C(=O)-, -C(=S)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(=N-CN)-, -C(=CH-NO<sub>2</sub>)-, -C[C(CN)<sub>2</sub>]-, -C(=CH-CN)-, -C(=NR<sup>11</sup>)- or -C(=N-S(=O)<sub>2</sub>R<sup>11a</sup>)-,

25 wherein R<sup>11</sup> is

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with  
30 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

5 aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

10

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino;

15

R<sup>11a</sup> is

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

20

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroarylamino,

25

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino,

30

Y is a valence bond, -O- or -N(R<sup>12</sup>)-,



wherein R<sup>12</sup> is

hydrogen,

5

C<sub>1-6</sub>-alkyl optionally substituted with

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, heteroaryl-amino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

10

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-amino or heteroaryl-amino,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which are optionally substituted with

15

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino,

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

20

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

A is a valence bond, C<sub>1-8</sub>-alkylene, C<sub>2-8</sub>-alkenylene, C<sub>2-8</sub>-alkynylene, C<sub>3-8</sub>-cyclo-alkylene or phenylene, or

25

when Y is -N(R<sup>12</sup>)-, A, together with R<sup>12</sup> and the nitrogen atom to which they are connected, may form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring system optionally containing one or more further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl, aryl,

30

heteroaryl, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroarylamino; and

Z is -R<sup>13</sup>, -OR<sup>13</sup>, -SR<sup>13</sup>, -NR<sup>13</sup>R<sup>14</sup>, -CHR<sup>13</sup>R<sup>14</sup>, -CR<sup>13</sup>R<sup>14</sup>R<sup>15</sup> or =CR<sup>13</sup>R<sup>14</sup>,

5

wherein R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> independently are

hydrogen,

10

C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl, which are optionally substituted with aryl, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, heteroaryl or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with

15

C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, hydroxy, amino, halogen, cyano or trifluoromethyl,

20

aryl, C<sub>3-15</sub>-cycloalkyl, aroyl or heteroaryl, which are optionally substituted with aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl, nitro, arylamino, heteroarylamino, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub>-alkylsulfonyl, sulfonylamino, arylthio, heteroarylthio, aryloxy, acylamino, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, C<sub>3-8</sub>-cycloalkanecarbonyl, hydroxy, amino, halogen, cyano or trifluoromethyl, where

25

R<sup>13</sup> and R<sup>14</sup> or R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup>, when they do not represent hydrogen, may be joined by one or more bridging linkers such as a valence bond, C<sub>1-4</sub>-alkylene, C<sub>2-4</sub>-alkenylene, -O-, -S-, -N(R<sup>16</sup>)-, -C(=O)-, -S(=O)-, -S(=O)<sub>2</sub>-, -C(R<sup>16</sup>R<sup>17</sup>)-, phenylene, biphenylene, -O-C<sub>1-4</sub>-alkylene, -S-C<sub>1-4</sub>-alkylene, -N(R<sup>16</sup>)-C<sub>1-4</sub>-alkylene,

30

-N=C<sub>1-4</sub>-alkylene, -O-C<sub>2-4</sub>-alkenylene, -S-C<sub>2-4</sub>-alkenylene or -N(R<sup>16</sup>)-C<sub>2-4</sub>-alkenylene, to form a mono-, bi- or polycyclic ring system,

wherein R<sup>16</sup> and R<sup>17</sup> independently are

5

hydrogen,

C<sub>1-6</sub>-alkyl optionally substituted with

10

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino, hetero-  
arylamino or C<sub>3-8</sub>-cycloalkyl, which are optionally substituted with  
C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen,  
cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, aryl-  
sulfonyl, arylamino or heteroarylamino,

15

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl, which  
are optionally substituted with  
C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano,  
trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, aryl-  
amino or heteroarylamino,

20

C<sub>1-6</sub>-alkylsulfonyl optionally substituted with

C<sub>3-8</sub>-cycloalkyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen,  
cyano, trifluoromethyl, aryl, heteroaryl, aroyl, heteroaroyl, arylsul-  
fonyl, arylamino or heteroarylamino, or

25

R<sup>16</sup> and R<sup>17</sup>, together with the nitrogen atom to which they are connected,  
form a 3 to 8 membered, saturated or unsaturated, heterocyclic ring op-  
tionally containing one or more further heteroatoms selected from nitro-  
gen, oxygen and sulfur, and optionally substituted with C<sub>1-6</sub>-alkyl,

30

C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, halogen, cyano, trifluoromethyl,

aryl, heteroaryl, aroyl, heteroaroyl, arylsulfonyl, arylamino or heteroaryl-amino;

as well as any optical or geometric isomer or tautomeric form thereof including mix-  
5 tures of these or a pharmaceutically acceptable salt thereof for the preparation of a  
medicament for the treatment of disorders related to the histamine H3 receptor.

35. Use of a compound as defined in claim 34 for the preparation of a medicament  
for the treatment and/or prevention of disorders and diseases wherein a histamine  
10 H3 antagonistic action is beneficial.

36. Use of a compound as defined in claim 34 for the preparation of a medicament  
for the reduction of weight.

15 37. Use of a compound as defined in claim 34 for the preparation of a medicament  
for the treatment and/or prevention of overweight or obesity.

38. Use of a compound as defined in claim 34 for the preparation of a medicament  
for the suppression of appetite or satiety induction.

20 39. Use of a compound as defined in claim 34 for the preparation of a medicament  
for the preparation of a medicament for the prevention and/or treatment of disorders  
and diseases related to overweight or obesity.

25 40. Use of a compound as defined in claim 34 for the preparation of a medicament  
for the prevention and/or treatment of eating disorders such as bulimia and binge  
eating.

30 41. Use of a compound as defined in claim 34 for the preparation of a medicament  
for use in the treatment of disorders related to the serotonin-3 receptor (5-HT3), such  
as for use in the treatment of emesis.

42. Use of a compound as defined in claim 34 for the preparation of a medicament for use in the treatment of disorders related to the vanilloid receptor, such as for use in the treatment of pain, neurogenic inflammation or obesity.

5

43. Use of a compound as defined in claim 34 for the preparation of a medicament for use in the treatment of disorders related to the alpha-2 adrenergic receptor, such as for use as a sleep inducing agent.

10

44. A method for the treatment of disorders related to the histamine H3 receptor the method comprising administering to a subject in need thereof an effective amount of a compound as defined in claim 34 or a pharmaceutical composition comprising the compound.

15

45. The method according to claim 44 wherein the effective amount of the compound as defined in claim 34 is in the range of from about 0.05 mg to about 2000 mg, preferably from about 0.1 mg to about 1000 mg and especially preferred from about 0.5 mg to about 500 mg per day.

20

46. Any novel feature or combination of features as described herein.

**ABSTRACT**

A novel class of substituted imidazole derivatives, methods for their preparation, pharmaceutical compositions comprising them and use thereof in the treatment of disorders related to the histamine H3 receptor. More particularly, the compounds  
5 possess histamine H3 receptor antagonistic activity and are thus useful in the treatment of disorders in which a histamine H3 receptor blockade is beneficial.